

**Anti-BMP6 Rabbit Monoclonal Antibody**  
**Catalog # ABO15569****Specification**

---

**Anti-BMP6 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, FC
Primary Accession	<a href="#">P22004</a>
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-BMP6 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications.  
This antibody reacts with Human, Mouse, Rat.

**Anti-BMP6 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 654

**Other Names**

Bone morphogenetic protein 6, BMP-6, VG-1-related protein, VG-1-R, VGR-1, BMP6, VGR

**Calculated MW**

42 kDa KDa

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>FC 1:50

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human BMP6

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-BMP6 Rabbit Monoclonal Antibody - Protein Information**

**Name** BMP6

## Synonyms VGR

### Function

Growth factor of the TGF-beta superfamily that plays essential roles in many developmental processes including cartilage and bone formation (PubMed:<a href="http://www.uniprot.org/citations/31019025" target="\_blank">31019025</a>). Also plays an important role in the regulation of HAMP/hepcidin expression and iron metabolism by acting as a ligand for hemojuvelin/HJV (PubMed:<a href="http://www.uniprot.org/citations/26582087" target="\_blank">26582087</a>). Also acts to promote expression of HAMP, potentially via the interaction with its receptor BMPR1A/ALK3 (PubMed:<a href="http://www.uniprot.org/citations/30097509" target="\_blank">30097509</a>, PubMed:<a href="http://www.uniprot.org/citations/31800957" target="\_blank">31800957</a>). Initiates the canonical BMP signaling cascade by associating with type I receptor ACVR1 and type II receptor ACVR2B (PubMed:<a href="http://www.uniprot.org/citations/18070108" target="\_blank">18070108</a>). In turn, ACVR1 propagates signal by phosphorylating SMAD1/5/8 that travel to the nucleus and act as activators and repressors of transcription of target. Can also signal through non-canonical pathway such as TAZ-Hippo signaling cascade to modulate VEGF signaling by regulating VEGFR2 expression (PubMed:<a href="http://www.uniprot.org/citations/33021694" target="\_blank">33021694</a>).

### Cellular Location

Secreted.

## Anti-BMP6 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-BMP6 Rabbit Monoclonal Antibody - Images




Figure 1. Western blot analysis of BMP6 using anti-BMP6 antibody (M06924-1). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,

Lane 2: human SiHa whole cell lysates,

Lane 3: human HEL whole cell lysates,

Lane 4: human 293T whole cell lysates,

Lane 5: rat PC-12 whole cell lysates,

Lane 6: rat RH35 whole cell lysates,

Lane 7: mouse HEPA1-6 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-BMP6 antigen affinity purified monoclonal antibody (Catalog # M06924-1) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:1000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for BMP6 at approximately 42 kDa. The expected band size for BMP6 is at 57 kDa.