

**Anti-Placental Protein 14 / PAEP Rabbit Monoclonal Antibody**  
**Catalog # ABO15590****Specification****Anti-Placental Protein 14 / PAEP Rabbit Monoclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P09466</a>
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-Placental Protein 14 / PAEP Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human.

**Anti-Placental Protein 14 / PAEP Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 5047

**Other Names**

Glycodelin, GD, Placental protein 14, PP14, Pregnancy-associated endometrial alpha-2 globulin, PAEG, PEG, Progesterone-associated endometrial protein, Progesterone-associated endometrial protein, Zona-binding inhibitory factor-1, ZIF-1, PAEP

**Calculated MW**

24 kDa KDa

**Application Details**

WB 1:500-1:2000

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human Placental Protein 14 / PAEP

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-Placental Protein 14 / PAEP Rabbit Monoclonal Antibody - Protein Information**

**Name** PAEP**Function**

Glycoprotein that regulates critical steps during fertilization and also has immunomodulatory effects. Four glycoforms, namely glycodelin-S, -A, -F and -C have been identified in reproductive tissues that differ in glycosylation and biological activity. Glycodelin-A has contraceptive and immunosuppressive activities (PubMed:<a href="http://www.uniprot.org/citations/7531163" target="\_blank">7531163</a>, PubMed:<a href="http://www.uniprot.org/citations/9918684" target="\_blank">9918684</a>). Glycodelin-C stimulates binding of spermatozoa to the zona pellucida (PubMed:<a href="http://www.uniprot.org/citations/17192260" target="\_blank">17192260</a>). Glycodelin-F inhibits spermatozoa-zona pellucida binding and significantly suppresses progesterone-induced acrosome reaction of spermatozoa (PubMed:<a href="http://www.uniprot.org/citations/12672671" target="\_blank">12672671</a>). Glycodelin-S in seminal plasma maintains the uncapacitated state of human spermatozoa (PubMed:<a href="http://www.uniprot.org/citations/15883155" target="\_blank">15883155</a>).

**Cellular Location**

Secreted

**Tissue Location**


This protein is, the main protein synthesized and secreted in the endometrium from mid-luteal phase of the menstrual cycle and during the first semester of pregnancy (PubMed:3667877) Glycodelin-A is expressed in amniotic fluid, endometrium/decidua and maternal serum (at protein level) (PubMed:3194393). Glycodelin-F is expressed in follicular fluid, luteinized granulosa cells and the oviduct (at protein level) (PubMed:12672671). Glycodelin-S is expressed in seminal plasma and seminal vesicles (at protein level) (PubMed:9239694). Glycodelin-C is detected in cumulus cells (at protein level), but cumulus cells do not synthesize Glycodelin-C but take up and convert glycodelin-A and -F via glycan remodeling (PubMed:17192260).

**Anti-Placental Protein 14 / PAEP Rabbit Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-Placental Protein 14 / PAEP Rabbit Monoclonal Antibody - Images**



Western blot analysis of Placental Protein 14 / PAEP expression in TPA-treated K562 cell lysate.