

Anti-CD97 Rabbit Monoclonal Antibody
Catalog # ABO15663**Specification**

Anti-CD97 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC, FC
Primary Accession	P48960
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-CD97 Rabbit Monoclonal Antibody . Tested in WB, IHC, Flow Cytometry applications. This antibody reacts with Human, Rat.

Anti-CD97 Rabbit Monoclonal Antibody - Additional Information

Gene ID 976

Other Names

Adhesion G protein-coupled receptor E5 {ECO:0000312|HGNC:HGNC:1711}, Leukocyte antigen CD97, CD97, Adhesion G protein-coupled receptor E5 subunit alpha, Adhesion G protein-coupled receptor E5 subunit beta, ADGRE5 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=1711)
HGNC:1711

Calculated MW

55 kDa, 28 kDa KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
FC 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human CD97

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-CD97 Rabbit Monoclonal Antibody - Protein Information

Name ADGRE5 ([HGNC:1711](#))

Function

Receptor potentially involved in both adhesion and signaling processes early after leukocyte activation. Plays an essential role in leukocyte migration.

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:Q9Z0M6}; Multi-pass membrane protein

Tissue Location


Broadly expressed, found on most hematopoietic cells, including activated lymphocytes, monocytes, macrophages, dendritic cells, and granulocytes. Expressed also abundantly by smooth muscle cells. Expressed in thyroid, colorectal, gastric, esophageal and pancreatic carcinomas too. Expression are increased under inflammatory conditions in the CNS of multiple sclerosis and in synovial tissue of patients with rheumatoid arthritis. Increased expression of CD97 in the synovium is accompanied by detectable levels of soluble CD97 in the synovial fluid

Anti-CD97 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CD97 Rabbit Monoclonal Antibody - Images



Western blot analysis of CD97 expression in human tonsil cell lysate.