

**Anti-Engrailed 1 Rabbit Monoclonal Antibody**  
**Catalog # ABO15803****Specification****Anti-Engrailed 1 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IP, FC
Primary Accession	<a href="#">Q05925</a>
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-Engrailed 1 Rabbit Monoclonal Antibody . Tested in WB, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

**Anti-Engrailed 1 Rabbit Monoclonal Antibody - Additional Information****Gene ID** 2019**Other Names**

Homeobox protein engrailed-1, Homeobox protein en-1, Hu-En-1, EN1

**Calculated MW**

55 kDa KDa

**Application Details**

WB 1:500-1:2000<br>IP 1:50<br>FC 1:100

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human Engrailed 1

**Purification**

Affinity-chromatography

**Storage**

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

**Anti-Engrailed 1 Rabbit Monoclonal Antibody - Protein Information****Name** EN1

**Function**

Required for proper formation of the apical ectodermal ridge and correct dorsal-ventral patterning in the limb.


**Cellular Location**

Nucleus.

**Anti-Engrailed 1 Rabbit Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-Engrailed 1 Rabbit Monoclonal Antibody - Images**

Western blot analysis of Engrailed 1 expression in 293T cell lysate.




Figure 1. Western blot analysis of EN1 using anti-EN1 antibody (M05395).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: rat brain tissue lysates,

Lane 2: mouse brain tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-EN1 antigen affinity purified monoclonal antibody (Catalog # M05395) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for EN1 at approximately 55 kDa. The expected band size for EN1 is at 40 kDa.