

Anti-PR3 Rabbit Monoclonal Antibody
Catalog # ABO15892**Specification**

Anti-PR3 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC
Primary Accession	P24158
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-PR3 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human.

Anti-PR3 Rabbit Monoclonal Antibody - Additional Information

Gene ID 5657

Other Names

Myeloblastin, 3.4.21.76, AGP7, C-ANCA antigen, Leukocyte proteinase 3, PR-3, PR3, Neutrophil proteinase 4, NP-4, P29, Wegener autoantigen, PRTN3, MBN

Calculated MW

27 kDa KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human PR3

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-PR3 Rabbit Monoclonal Antibody - Protein Information

Name PRTN3

Synonyms MBN

Function

Serine protease that degrades elastin, fibronectin, laminin, vitronectin, and collagen types I, III, and IV (in vitro) (PubMed:2033050, PubMed:28240246, PubMed:3198760). By cleaving and activating receptor F2RL1/PAR-2, enhances endothelial cell barrier function and thus vascular integrity during neutrophil transendothelial migration (PubMed:23202369). Plays a role in neutrophil transendothelial migration, probably when associated with CD177 (PubMed:22266279). Triggers inflammatory processes in neutrophils by interacting with ADGRG3 upstream of F2RL1/PAR2 activation (PubMed:36302784).

Cellular Location

Cytoplasmic granule. Secreted. Cell membrane; Peripheral membrane protein; Extracellular side. Membrane raft; Peripheral membrane protein; Extracellular side. Note=Localizes predominantly to azurophil granules (primary secretory granules) in neutrophils (PubMed:18462208, PubMed:2033050, PubMed:3198760, PubMed:7897245) Secreted upon neutrophil stimulation by TNF-alpha, lipopolysaccharide (LPS), fMLP and CXCL8/IL8 or during neutrophil transmigration (PubMed:22266279, PubMed:28240246). Following secretion tethered to the cell membrane by CD177 (PubMed:18462208, PubMed:22266279)

Tissue Location


Expressed in polymorphonuclear leukocytes (at protein level) (PubMed:2033050, PubMed:3198760, PubMed:7897245) Expressed in neutrophils (at protein level) (PubMed:17244676, PubMed:18462208, PubMed:21193407, PubMed:22266279, PubMed:28240246) Expressed in differentiating neutrophils (PubMed:18462208)


Anti-PR3 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-PR3 Rabbit Monoclonal Antibody - Images





Western blot analysis of PR3 expression in HL60 cell lysate.



All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.