

Anti-FRA1 Rabbit Monoclonal Antibody
Catalog # ABO15894**Specification**

Anti-FRA1 Rabbit Monoclonal Antibody - Product Information

Application	WB, IF, ICC
Primary Accession	P15407
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-FRA1 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF applications. This antibody reacts with Human.

Anti-FRA1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 8061

Other Names

Fos-related antigen 1, FRA-1, FOSL1, FRA1

Calculated MW

39 kDa KDa

Application Details

WB 1:500-1:2000
ICC/IF 1:50-1:200

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human FRA1

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-FRA1 Rabbit Monoclonal Antibody - Protein Information

Name FOSL1

Synonyms FRA1

Cellular Location

Nucleus {ECO:0000250|UniProtKB:P51145}.

Anti-FRA1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-FRA1 Rabbit Monoclonal Antibody - Images




Figure 1. Western blot analysis of FRA1 using anti-FRA1 antibody (M03927-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human MOLT-4 whole cell lysates,

Lane 2: human A431 whole cell lysates,

Lane 3: human U251 whole cell lysates,

Lane 4: human A549 whole cell lysates,

Lane 5: human SIHA whole cell lysates,

Lane 6: human U20S whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-FRA1 antigen affinity purified monoclonal antibody (Catalog # M03927-1) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for FRA1 at

approximately 39 kDa. The expected band size for FRA1 is at 29 kDa.