

Anti-Inhibin beta B Rabbit Monoclonal Antibody

Catalog # ABO15919

Specification

Anti-Inhibin beta B Rabbit Monoclonal Antibody - Product Information

Application WB
Primary Accession P09529
Host Rabbit Isotype IgG

Reactivity Rat, Human, Mouse

Clonality Monoclonal Format Liquid

Description

Anti-Inhibin beta B Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Anti-Inhibin beta B Rabbit Monoclonal Antibody - Additional Information

Gene ID 3625

Other Names

Inhibin beta B chain, Activin beta-B chain, INHBB

Calculated MW 52 kDa KDa

Application Details WB 1:500-1:2000

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human Inhibin beta B

Purification

Affinity-chromatography

Storage Store at -20°C for one year. For short term


storage and frequent use, store at 4°C for

up to one month. Avoid repeated

freeze-thaw cycles.

Anti-Inhibin beta B Rabbit Monoclonal Antibody - Protein Information

Name INHBB





Function

Inhibins and activins inhibit and activate, respectively, the secretion of follitropin by the pituitary gland. Inhibins/activins are involved in regulating a number of diverse functions such as hypothalamic and pituitary hormone secretion, gonadal hormone secretion, germ cell development and maturation, erythroid differentiation, insulin secretion, nerve cell survival, embryonic axial development or bone growth, depending on their subunit composition. Inhibins appear to oppose the functions of activins. Inhibin B is a dimer of alpha and beta-B that plays a crucial role in the regulation of the reproductive system by inhibiting the secretion of follicle-stimulating hormone (FSH) from the anterior pituitary gland. Thereby, maintains reproductive homeostasis in both males and females. Acts as a more potent suppressor of FSH release than inhibin A (By similarity). Functions as competitive receptor antagonist binding activin type II receptors with high affinity in the presence of the TGF-beta type III coreceptor/TGFBR3L (By similarity).


Cellular Location Secreted.

Anti-Inhibin beta B Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-Inhibin beta B Rabbit Monoclonal Antibody - Images



Western blot analysis of Inhibin beta B expression in Mouse brain lysate.