

Anti-eIF3e Rabbit Monoclonal Antibody
Catalog # ABO15937**Specification****Anti-eIF3e Rabbit Monoclonal Antibody - Product Information**

Application	WB, IF, ICC
Primary Accession	P60228
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-eIF3e Rabbit Monoclonal Antibody . Tested in WB, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

Anti-eIF3e Rabbit Monoclonal Antibody - Additional Information

Gene ID 3646

Other Names

Eukaryotic translation initiation factor 3 subunit E {ECO:0000255|HAMAP-Rule:MF_03004}, eIF3e {ECO:0000255|HAMAP-Rule:MF_03004}, Eukaryotic translation initiation factor 3 subunit 6 {ECO:0000255|HAMAP-Rule:MF_03004}, Viral integration site protein INT-6 homolog, eIF-3 p48 {ECO:0000255|HAMAP-Rule:MF_03004}, EIF3E {ECO:0000255|HAMAP-Rule:MF_03004}

Calculated MW

52 kDa KDa

Application Details

WB 1:500-1:2000
ICC/IF 1:50-1:200

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human eIF3e

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-eIF3e Rabbit Monoclonal Antibody - Protein Information

Name EIF3E {ECO:0000255|HAMAP-Rule:MF_03004}

Function

Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed:17581632, PubMed:25849773, PubMed:27462815). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl- tRNA_i and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed:17581632). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed:25849773). Required for nonsense-mediated mRNA decay (NMD); may act in conjunction with UPF2 to divert mRNAs from translation to the NMD pathway (PubMed:17468741). May interact with MCM7 and EPAS1 and regulate the proteasome-mediated degradation of these proteins (PubMed:17310990, PubMed:17324924).

Cellular Location

Cytoplasm. Nucleus, PML body.

Tissue Location


Ubiquitously expressed. Expressed at highest levels in appendix, lymph, pancreas, skeletal muscle, spleen and thymus


Anti-eIF3e Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-eIF3e Rabbit Monoclonal Antibody - Images





Western blot analysis of eIF3e expression in (1) 293T cell lysate; (2) Jurkat cell lysate.