

Anti-BAT3 Rabbit Monoclonal Antibody
Catalog # ABO15978**Specification****Anti-BAT3 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, FC
Primary Accession	P46379
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-BAT3 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications.
This antibody reacts with Human, Mouse, Rat.

Anti-BAT3 Rabbit Monoclonal Antibody - Additional Information

Gene ID 7917

Other Names

Large proline-rich protein BAG6, BAG family molecular chaperone regulator 6, BCL2-associated athanogene 6 {ECO:0000312|HGNC:HGNC:13919}, BAG-6, HLA-B-associated transcript 3, Protein G3, Protein Scythe, BAG6 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=13919)
HGNC:13919

Calculated MW

150 kDa KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
FC 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human BAT3

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-BAT3 Rabbit Monoclonal Antibody - Protein Information

Name BAG6 ([HGNC:13919](#))

Function

ATP-independent molecular chaperone preventing the aggregation of misfolded and hydrophobic patches-containing proteins (PubMed:[21636303](http://www.uniprot.org/citations/21636303)). Functions as part of a cytosolic protein quality control complex, the BAG6/BAT3 complex, which maintains these client proteins in a soluble state and participates in their proper delivery to the endoplasmic reticulum or alternatively can promote their sorting to the proteasome where they undergo degradation (PubMed:[20516149](http://www.uniprot.org/citations/20516149), PubMed:[21636303](http://www.uniprot.org/citations/21636303), PubMed:[21743475](http://www.uniprot.org/citations/21743475), PubMed:[28104892](http://www.uniprot.org/citations/28104892)). The BAG6/BAT3 complex is involved in the post-translational delivery of tail-anchored/type II transmembrane proteins to the endoplasmic reticulum membrane. Recruited to ribosomes, it interacts with the transmembrane region of newly synthesized tail-anchored proteins and together with SGTA and ASNA1 mediates their delivery to the endoplasmic reticulum (PubMed:[20516149](http://www.uniprot.org/citations/20516149), PubMed:[20676083](http://www.uniprot.org/citations/20676083), PubMed:[25535373](http://www.uniprot.org/citations/25535373), PubMed:[28104892](http://www.uniprot.org/citations/28104892)). Client proteins that cannot be properly delivered to the endoplasmic reticulum are ubiquitinated by RNF126, an E3 ubiquitin-protein ligase associated with BAG6 and are sorted to the proteasome (PubMed:[24981174](http://www.uniprot.org/citations/24981174), PubMed:[27193484](http://www.uniprot.org/citations/27193484), PubMed:[28104892](http://www.uniprot.org/citations/28104892)). SGTA which prevents the recruitment of RNF126 to BAG6 may negatively regulate the ubiquitination and the proteasomal degradation of client proteins (PubMed:[23129660](http://www.uniprot.org/citations/23129660), PubMed:[25179605](http://www.uniprot.org/citations/25179605), PubMed:[27193484](http://www.uniprot.org/citations/27193484)). Similarly, the BAG6/BAT3 complex also functions as a sorting platform for proteins of the secretory pathway that are mislocalized to the cytosol either delivering them to the proteasome for degradation or to the endoplasmic reticulum (PubMed:[21743475](http://www.uniprot.org/citations/21743475)). The BAG6/BAT3 complex also plays a role in the endoplasmic reticulum-associated degradation (ERAD), a quality control mechanism that eliminates unwanted proteins of the endoplasmic reticulum through their retrotranslocation to the cytosol and their targeting to the proteasome. It maintains these retrotranslocated proteins in an unfolded yet soluble state condition in the cytosol to ensure their proper delivery to the proteasome (PubMed:[21636303](http://www.uniprot.org/citations/21636303)). BAG6 is also required for selective ubiquitin-mediated degradation of defective nascent chain polypeptides by the proteasome. In this context, it may participate in the production of antigenic peptides and play a role in antigen presentation in immune response (By similarity). BAG6 is also involved in endoplasmic reticulum stress-induced pre-emptive quality control, a mechanism that selectively attenuates the translocation of newly synthesized proteins into the endoplasmic reticulum and reroutes them to the cytosol for proteasomal degradation. BAG6 may ensure the proper degradation of these proteins and thereby protects the endoplasmic reticulum from protein overload upon stress (PubMed:[26565908](http://www.uniprot.org/citations/26565908)). By inhibiting the polyubiquitination and subsequent proteasomal degradation of HSPA2 it may also play a role in the assembly of the synaptonemal complex during spermatogenesis (By similarity). Also positively regulates apoptosis by interacting with and stabilizing the proapoptotic factor AIFM1 (By similarity). By controlling the steady-state expression of the IGF1R receptor, indirectly regulates the insulin-like growth factor receptor signaling pathway (PubMed:[26692333](http://www.uniprot.org/citations/26692333)).

Cellular Location

Cytoplasm, cytosol. Nucleus. Secreted, extracellular exosome Note=Normally localized in cytosol and nucleus, it can also be released extracellularly, in exosomes, by tumor and myeloid dendritic cells (PubMed:18055229, PubMed:18852879). Cytoplasmic retention is due to interaction with GET4 (PubMed:29042515).


Tissue Location

Expressed by immature dendritic cells (at protein level).

Anti-BAT3 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-BAT3 Rabbit Monoclonal Antibody - Images

Western blot analysis of BAT3 expression in A431 cell lysate.