

Anti-PEX19 Rabbit Monoclonal Antibody
Catalog # ABO15979**Specification****Anti-PEX19 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IF, ICC, IP, FC
Primary Accession	P40855
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-PEX19 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Rat.

Anti-PEX19 Rabbit Monoclonal Antibody - Additional Information**Gene ID 5824****Other Names**

Peroxisomal biogenesis factor 19, 33 kDa housekeeping protein, Peroxin-19, Peroxisomal farnesylated protein, PEX19 (HGNC:9713), HK33, PXF

Calculated MW

35 kDa KDa

Application Details

WB 1:500-1:2000
ICC/IF 1:50-1:200
IP 1:50
FC 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human PEX19

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-PEX19 Rabbit Monoclonal Antibody - Protein Information

Name PEX19 ([HGNC:9713](#))

Synonyms HK33, PXF

Function

Necessary for early peroxisomal biogenesis. Acts both as a cytosolic chaperone and as an import receptor for peroxisomal membrane proteins (PMPs). Binds and stabilizes newly synthesized PMPs in the cytoplasm by interacting with their hydrophobic membrane-spanning domains, and targets them to the peroxisome membrane by binding to the integral membrane protein PEX3. Excludes CDKN2A from the nucleus and prevents its interaction with MDM2, which results in active degradation of TP53.

Cellular Location

Cytoplasm. Peroxisome membrane; Lipid-anchor; Cytoplasmic side. Note=Mainly cytoplasmic. Some fraction membrane-associated to the outer surface of peroxisomes.

Tissue Location


Ubiquitously expressed. Isoform 1 is strongly predominant in all tissues except in utero where isoform 2 is the main form.

Anti-PEX19 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-PEX19 Rabbit Monoclonal Antibody - Images



Western blot analysis of PEX19 expression in Jurkat cell lysate.