

# **Anti-PDLIM1 Rabbit Monoclonal Antibody**

**Catalog # ABO15989** 

## **Specification**

### **Anti-PDLIM1 Rabbit Monoclonal Antibody - Product Information**

Application WB, IF, ICC, FC

Primary Accession

Host
Isotype

O00151

Rabbit
IgG

Reactivity Rat, Human, Mouse

Clonality Monoclonal Format Liquid

**Description** 

Anti-PDLIM1 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

## **Anti-PDLIM1 Rabbit Monoclonal Antibody - Additional Information**

#### **Gene ID 9124**

#### **Other Names**

PDZ and LIM domain protein 1, C-terminal LIM domain protein 1, Elfin, LIM domain protein CLP-36, PDLIM1, CLIM1, CLP36

# **Calculated MW**

36 kDa KDa

### **Application Details**

WB 1:500-1:2000<br>ICC/IF 1:50-1:200<br>FC 1:50</br>

#### **Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

#### **Immunogen**

A synthesized peptide derived from human PDLIM1

# **Purification**

Affinity-chromatography

Storage Store at -20°C for one year. For short term

storage and frequent use, store at 4°C for

up to one month. Avoid repeated

freeze-thaw cycles.

### Anti-PDLIM1 Rabbit Monoclonal Antibody - Protein Information

## Name PDLIM1



### Synonyms CLIM1, CLP36

#### **Function**

Cytoskeletal protein that may act as an adapter that brings other proteins (like kinases) to the cytoskeleton (PubMed:<a href="http://www.uniprot.org/citations/10861853" target="\_blank">10861853</a>). Involved in assembly, disassembly and directioning of stress fibers in fibroblasts. Required for the localization of ACTN1 and PALLD to stress fibers. Required for cell migration and in maintaining cell polarity of fibroblasts (By similarity).

### **Cellular Location**

Cytoplasm. Cytoplasm, cytoskeleton. Cytoplasm, myofibril, sarcomere, Z line Note=Associates with actin stress fibers

#### **Tissue Location**


Strongly expressed in the heart and skeletal muscle, moderately expressed in the spleen, small intestine, colon, placenta, and lung. A lower level expression is seen in liver, thymus, kidney, prostate and pancreas and is not found in the brain, testis, ovary, and peripheral blood leukocytes

### **Anti-PDLIM1 Rabbit Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### **Anti-PDLIM1 Rabbit Monoclonal Antibody - Images**



Western blot analysis of PDLIM1 expression in 293T cell lysate.