

Anti-Smg1 Rabbit Monoclonal Antibody
Catalog # ABO15996**Specification****Anti-Smg1 Rabbit Monoclonal Antibody - Product Information**

Application	WB
Primary Accession	Q96Q15
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-Smg1 Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Anti-Smg1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 23049

Other Names

Serine/threonine-protein kinase SMG1, SMG-1, hSMG-1, 2.7.11.1, Lambda/iota protein kinase C-interacting protein, Lambda-interacting protein, Nonsense mediated mRNA decay-associated PI3K-related kinase SMG1 {ECO:0000312|HGNC:HGNC:30045}, SMG1 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=30045)

Calculated MW

410 kDa KDa

Application Details

WB 1:500-1:2000

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human Smg1

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-Smg1 Rabbit Monoclonal Antibody - Protein Information

Name SMG1 ([HGNC:30045](#))

Function

Serine/threonine protein kinase involved in both mRNA surveillance and genotoxic stress response pathways. Recognizes the substrate consensus sequence [ST]-Q. Plays a central role in nonsense-mediated decay (NMD) of mRNAs containing premature stop codons by phosphorylating UPF1/RENT1. Recruited by release factors to stalled ribosomes together with SMG8 and SMG9 (forming the SMG1C protein kinase complex), and UPF1 to form the transient SURF (SMG1-UPF1-eRF1-eRF3) complex. In EJC-dependent NMD, the SURF complex associates with the exon junction complex (EJC) through UPF2 and allows the formation of an UPF1-UPF2-UPF3 surveillance complex which is believed to activate NMD. Also acts as a genotoxic stress-activated protein kinase that displays some functional overlap with ATM. Can phosphorylate p53/TP53 and is required for optimal p53/TP53 activation after cellular exposure to genotoxic stress. Its depletion leads to spontaneous DNA damage and increased sensitivity to ionizing radiation (IR). May activate PRKCI but not PRKCZ.

Cellular Location

Nucleus. Cytoplasm. Note=Present in the chromatoid body {ECO:0000250|UniProtKB:Q8BKX6}

Tissue Location


Widely expressed, with highest level in heart and skeletal muscle. Expressed in placenta, brain, lung and spleen, but not in liver.


Anti-Smg1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Smg1 Rabbit Monoclonal Antibody - Images





Western blot analysis of Smg1 expression in Saos2 cell lysate.