

**Anti-MLKL Rabbit Monoclonal Antibody**  
**Catalog # ABO16058****Specification**

---

**Anti-MLKL Rabbit Monoclonal Antibody - Product Information**

Application	WB, IP
Primary Accession	<a href="#">Q8NB16</a>
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-MLKL Rabbit Monoclonal Antibody . Tested in WB, IP applications. This antibody reacts with Human, Mouse, Rat.

**Anti-MLKL Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 197259

**Other Names**

Mixed lineage kinase domain-like protein, hMLKL, MLKL {ECO:0000303|PubMed:22265413, ECO:0000312|HGNC:HGNC:26617}

**Calculated MW**

54 kDa KDa

**Application Details**

WB 1:500-1:2000<br>IP 1:50

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from MLKL

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-MLKL Rabbit Monoclonal Antibody - Protein Information**

**Name** MLKL {ECO:0000303|PubMed:22265413, ECO:0000312|HGNC:HGNC:26617}

## Function

Pseudokinase that plays a key role in TNF-induced necroptosis, a programmed cell death process (PubMed:<a href="http://www.uniprot.org/citations/22265413" target="\_blank">22265413</a>, PubMed:<a href="http://www.uniprot.org/citations/22265414" target="\_blank">22265414</a>, PubMed:<a href="http://www.uniprot.org/citations/22421439" target="\_blank">22421439</a>, PubMed:<a href="http://www.uniprot.org/citations/24316671" target="\_blank">24316671</a>). Does not have protein kinase activity (PubMed:<a href="http://www.uniprot.org/citations/22265413" target="\_blank">22265413</a>, PubMed:<a href="http://www.uniprot.org/citations/22265414" target="\_blank">22265414</a>, PubMed:<a href="http://www.uniprot.org/citations/22421439" target="\_blank">22421439</a>, PubMed:<a href="http://www.uniprot.org/citations/24316671" target="\_blank">24316671</a>). Activated following phosphorylation by RIPK3, leading to homotrimerization, localization to the plasma membrane and execution of programmed necrosis characterized by calcium influx and plasma membrane damage (PubMed:<a href="http://www.uniprot.org/citations/22265413" target="\_blank">22265413</a>, PubMed:<a href="http://www.uniprot.org/citations/22265414" target="\_blank">22265414</a>, PubMed:<a href="http://www.uniprot.org/citations/22421439" target="\_blank">22421439</a>, PubMed:<a href="http://www.uniprot.org/citations/24316671" target="\_blank">24316671</a>). In addition to TNF-induced necroptosis, necroptosis can also take place in the nucleus in response to orthomyxoviruses infection: following activation by ZBP1, MLKL is phosphorylated by RIPK3 in the nucleus, triggering disruption of the nuclear envelope and leakage of cellular DNA into the cytosol. following ZBP1 activation, which senses double-stranded Z-RNA structures, nuclear RIPK3 catalyzes phosphorylation and activation of MLKL, promoting disruption of the nuclear envelope and leakage of cellular DNA into the cytosol (By similarity). Binds to highly phosphorylated inositol phosphates such as inositolhexakisphosphate (InsP6) which is essential for its necroptotic function (PubMed:<a href="http://www.uniprot.org/citations/29883610" target="\_blank">29883610</a>).

## Cellular Location


Cytoplasm. Cell membrane Nucleus {ECO:0000250|UniProtKB:Q9D2Y4}. Note=Localizes to the cytoplasm and translocates to the plasma membrane on necroptosis induction (PubMed:24316671). Localizes to the nucleus in response to orthomyxoviruses infection (By similarity) {ECO:0000250|UniProtKB:Q9D2Y4, ECO:0000269|PubMed:24316671}

## Anti-MLKL Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-MLKL Rabbit Monoclonal Antibody - Images






Figure 1. Western blot analysis of MLKL using anti-MLKL antibody (M00535-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HeLa whole cell lysates,

Lane 2: human PC-3 whole cell lysates,

Lane 3: human A549 whole cell lysates,

Lane 4: human MCF-7 whole cell lysates,

Lane 5: rat testis tissue lysates,

Lane 6: rat liver tissue lysates,

Lane 7: mouse testis tissue lysates,

Lane 8: mouse liver tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MLKL antigen affinity purified monoclonal antibody (Catalog # M00535-1) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for MLKL at approximately 54 kDa. The expected band size for MLKL is at 54 kDa.