

**Anti-USP39 Rabbit Monoclonal Antibody**  
**Catalog # ABO16153****Specification****Anti-USP39 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC
Primary Accession	<a href="#">Q53GS9</a>
Host	Rabbit
Isotype	IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-USP39 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse.

**Anti-USP39 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 10713

**Other Names**

Ubiquitin carboxyl-terminal hydrolase 39, 3.4.19.12, SAD1 homolog, U4/U6.U5 tri-snRNP-associated 65 kDa protein, USP39 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=20071](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=20071))  
HGNC:20071

**Calculated MW**

65 kDa KDa

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human USP39

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-USP39 Rabbit Monoclonal Antibody - Protein Information**

**Name** USP39 ([HGNC:20071](#))

### Function

Deubiquitinating enzyme that plays a role in many cellular processes including cellular antiviral response, epithelial morphogenesis, DNA repair or B-cell development (PubMed:[33127822](http://www.uniprot.org/citations/33127822), PubMed:[34614178](http://www.uniprot.org/citations/34614178)). Plays a role in pre-mRNA splicing as a component of the U4/U6-U5 tri-snRNP, one of the building blocks of the precatalytic spliceosome (PubMed:[11350945](http://www.uniprot.org/citations/11350945), PubMed:[26912367](http://www.uniprot.org/citations/26912367)). Specifically regulates immunoglobulin gene rearrangement in a spliceosome-dependent manner, which involves modulating chromatin interactions at the Igh locus and therefore plays an essential role in B-cell development (By similarity). Regulates AURKB mRNA levels, and thereby plays a role in cytokinesis and in the spindle checkpoint (PubMed:[18728397](http://www.uniprot.org/citations/18728397)). Regulates apoptosis and G2/M cell cycle checkpoint in response to DNA damage by deubiquitinating and stabilizing CHK2 (PubMed:[30771428](http://www.uniprot.org/citations/30771428)). Also plays an important role in DNA repair by controlling the recruitment of XRCC4/LIG4 to DNA double-strand breaks for non-homologous end-joining repair (PubMed:[34614178](http://www.uniprot.org/citations/34614178)). Participates in antiviral activity by affecting the type I IFN signaling by stabilizing STAT1 and decreasing its 'Lys-6'-linked ubiquitination (PubMed:[33127822](http://www.uniprot.org/citations/33127822)). Contributes to non-canonical Wnt signaling during epidermal differentiation (By similarity). Acts as a negative regulator NF-kappa-B activation through deubiquitination of 'Lys-48'-linked ubiquitination of NFKBIA (PubMed:[36651806](http://www.uniprot.org/citations/36651806)).

### Cellular Location


Nucleus


### Anti-USP39 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-USP39 Rabbit Monoclonal Antibody - Images





Western blot analysis of USP39 expression in 293T cell lysate.