

**Anti-Phospho-MCM2 (S27) Rabbit Monoclonal Antibody**  
**Catalog # ABO16329****Specification****Anti-Phospho-MCM2 (S27) Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, IP
Primary Accession	<a href="#">P49736</a>
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-Phospho-MCM2 (S27) Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human, Mouse, Rat.

**Anti-Phospho-MCM2 (S27) Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 4171

**Other Names**

DNA replication licensing factor MCM2, 3.6.4.12, Minichromosome maintenance protein 2 homolog, Nuclear protein BM28, MCM2 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=6944](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=6944))  
HGNC:6944

**Calculated MW**

125 kDa KDa

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>IP 1:50

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human Phospho-MCM2 (S27)

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-Phospho-MCM2 (S27) Rabbit Monoclonal Antibody - Protein Information**

**Name** MCM2 ([HGNC:6944](#))

### Function

Acts as a component of the MCM2-7 complex (MCM complex) which is the replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. Core component of CDC45-MCM-GINS (CMG) helicase, the molecular machine that unwinds template DNA during replication, and around which the replisome is built (PubMed:<a href="http://www.uniprot.org/citations/32453425" target="\_blank">32453425</a>, PubMed:<a href="http://www.uniprot.org/citations/34694004" target="\_blank">34694004</a>, PubMed:<a href="http://www.uniprot.org/citations/34700328" target="\_blank">34700328</a>, PubMed:<a href="http://www.uniprot.org/citations/35585232" target="\_blank">35585232</a>). The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity (PubMed:<a href="http://www.uniprot.org/citations/32453425" target="\_blank">32453425</a>). Required for the entry in S phase and for cell division (PubMed:<a href="http://www.uniprot.org/citations/8175912" target="\_blank">8175912</a>). Plays a role in terminally differentiated hair cells development of the cochlea and induces cells apoptosis (PubMed:<a href="http://www.uniprot.org/citations/26196677" target="\_blank">26196677</a>).

### Cellular Location


Nucleus. Chromosome. Note=Associated with chromatin before the formation of nuclei and detaches from it as DNA replication progresses. {ECO:0000250|UniProtKB:P55861}


### Anti-Phospho-MCM2 (S27) Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-Phospho-MCM2 (S27) Rabbit Monoclonal Antibody - Images





Western blot analysis of Phospho-MCM2 (S27) expression in (1) HeLa cell lysate; (2) HeLa cell treated with alkaline phosphatase lysate.