

Anti-Importin 9 / RANBP9 Rabbit Monoclonal Antibody
Catalog # ABO16354**Specification****Anti-Importin 9 / RANBP9 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC
Primary Accession	Q96P70
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-Importin 9 / RANBP9 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human.

Anti-Importin 9 / RANBP9 Rabbit Monoclonal Antibody - Additional Information

Gene ID 55705

Other Names

Importin-9, Imp9, Ran-binding protein 9, RanBP9, IPO9 {ECO:0000303|PubMed:30855230, ECO:0000312|HGNC:HGNC:19425}

Calculated MW

140 kDa KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human Importin 9 / RANBP9

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-Importin 9 / RANBP9 Rabbit Monoclonal Antibody - Protein Information

Name IPO9 {ECO:0000303|PubMed:30855230, ECO:0000312|HGNC:HGNC:19425}

Function

Nuclear transport receptor that mediates nuclear import of proteins, such as histones, proteasome and actin (PubMed:11823430, PubMed:30855230, PubMed:34711951). Serves as receptor for nuclear localization signals (NLS) in cargo substrates (PubMed:11823430). Is thought to mediate docking of the importin/substrate complex to the nuclear pore complex (NPC) through binding to nucleoporin and the complex is subsequently translocated through the pore by an energy requiring, Ran-dependent mechanism (PubMed:11823430). At the nucleoplasmic side of the NPC, Ran binds to the importin, the importin/substrate complex dissociates and importin is re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran (PubMed:11823430). The directionality of nuclear import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus (PubMed:11823430). Mediates the import of pre-assembled proteasomes into the nucleus; AKIRIN2 acts as a molecular bridge between IPO9 and the proteasome complex (PubMed:11823430, PubMed:34711951). Mediates the nuclear import of histones H2A, H2B, H4 and H4 (PubMed:11823430, PubMed:30855230). In addition to nuclear import, also acts as a chaperone for histones by preventing inappropriate non-nucleosomal interactions (PubMed:30855230). Mediates the nuclear import of actin (By similarity).

Cellular Location


Cytoplasm. Nucleus


Anti-Importin 9 / RANBP9 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Importin 9 / RANBP9 Rabbit Monoclonal Antibody - Images





Western blot analysis of Importin 9 / RANBP9 expression in HeLa cell lysate.



All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.