

**Anti-GNA13 Rabbit Monoclonal Antibody**  
**Catalog # ABO16364****Specification**

---

**Anti-GNA13 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">Q14344</a>
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-GNA13 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human.

**Anti-GNA13 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 10672

**Other Names**

Guanine nucleotide-binding protein subunit alpha-13, G alpha-13, G-protein subunit alpha-13, GNA13

**Calculated MW**

44 kDa KDa

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human GNA13

**Purification**

Affinity-chromatography

Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-GNA13 Rabbit Monoclonal Antibody - Protein Information**

**Name** GNA13

### Function

Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems (PubMed:<a href="http://www.uniprot.org/citations/15240885" target="\_blank">15240885</a>, PubMed:<a href="http://www.uniprot.org/citations/16705036" target="\_blank">16705036</a>, PubMed:<a href="http://www.uniprot.org/citations/16787920" target="\_blank">16787920</a>, PubMed:<a href="http://www.uniprot.org/citations/27084452" target="\_blank">27084452</a>). Activates effector molecule RhoA by binding and activating RhoGEFs (ARHGEF1/p115RhoGEF, ARHGEF11/PDZ-RhoGEF and ARHGEF12/LARG) (PubMed:<a href="http://www.uniprot.org/citations/12515866" target="\_blank">12515866</a>, PubMed:<a href="http://www.uniprot.org/citations/15240885" target="\_blank">15240885</a>). GNA13-dependent Rho signaling subsequently regulates transcription factor AP-1 (activating protein-1) (By similarity). Promotes tumor cell invasion and metastasis by activating RhoA/ROCK signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/16705036" target="\_blank">16705036</a>, PubMed:<a href="http://www.uniprot.org/citations/16787920" target="\_blank">16787920</a>, PubMed:<a href="http://www.uniprot.org/citations/27084452" target="\_blank">27084452</a>). Inhibits CDH1-mediated cell adhesion in a process independent from Rho activation (PubMed:<a href="http://www.uniprot.org/citations/11976333" target="\_blank">11976333</a>). In lymphoid follicles, transmits P2RY8- and S1PR2-dependent signals that lead to inhibition of germinal center (GC) B cell growth and migration outside the GC niche.

### Cellular Location

Cell membrane; Lipid-anchor. Melanosome. Cytoplasm. Nucleus Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065). Detected in the cytoplasm of Leydig cells and in the seminiferous epithelium, including differentiating cells from the spermatogonia to mature spermatozoa stages (PubMed:18703424). In round spermatids, also present in the nuclei (PubMed:18703424).

### Tissue Location


Expressed in testis, including in Leydig cells and in the seminiferous epithelium, in differentiating cells from the spermatogonia to mature spermatozoa stages and round spermatids (at protein level). Expressed in 99.2% of spermatozoa from healthy individuals, but only in 28.6% of macrocephalic spermatozoa from infertile patients (at protein level).


## Anti-GNA13 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-GNA13 Rabbit Monoclonal Antibody - Images





Western blot analysis of GNA13 expression in HepG2 cell lysate.