

Anti-RNF14 Rabbit Monoclonal Antibody
Catalog # ABO16374**Specification**

Anti-RNF14 Rabbit Monoclonal Antibody - Product Information

Application	WB
Primary Accession	Q9UBS8
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-RNF14 Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Anti-RNF14 Rabbit Monoclonal Antibody - Additional Information

Gene ID 9604

Other Names

E3 ubiquitin-protein ligase RNF14, 2.3.2.31, Androgen receptor-associated protein 54, HFB30, RING finger protein 14, RNF14 {ECO:0000303|PubMed:36638793, ECO:0000312|HGNC:HGNC:10058}

Calculated MW

54 kDa KDa

Application Details

WB 1:500-1:2000

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human RNF14

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-RNF14 Rabbit Monoclonal Antibody - Protein Information

Name RNF14 {ECO:0000303|PubMed:36638793, ECO:0000312|HGNC:HGNC:10058}

Function

E3 ubiquitin-protein ligase that plays a key role in the RNF14-RNF25 translation quality control pathway, a pathway that takes place when a ribosome has stalled during translation, and which promotes ubiquitination and degradation of translation factors on stalled ribosomes (PubMed:36638793, PubMed:37651229, PubMed:37951215, PubMed:37951216). Recruited to stalled ribosomes by the ribosome collision sensor GCN1 and mediates 'Lys-6'-linked ubiquitination of target proteins, leading to their degradation (PubMed:36638793, PubMed:37651229, PubMed:37951215, PubMed:37951216). Mediates ubiquitination of EEF1A1/eEF1A and ETF1/eRF1 translation factors on stalled ribosomes, leading to their degradation (PubMed:36638793, PubMed:37651229). Also catalyzes ubiquitination of ribosomal proteins RPL0, RPL1, RPL12, RPS13 and RPS17 (PubMed:36638793). Specifically required to resolve RNA-protein cross-links caused by reactive aldehydes, which trigger translation stress by stalling ribosomes: acts by catalyzing 'Lys-6'-linked ubiquitination of RNA-protein cross-links, leading to their removal by the ATP-dependent unfoldase VCP and subsequent degradation by the proteasome (PubMed:37951215, PubMed:37951216). Independently of its function in the response to stalled ribosomes, acts as a regulator of transcription in Wnt signaling via its interaction with TCF transcription factors (TCF7/TCF1, TCF7L1/TCF3 and TCF7L2/TCF4) (PubMed:23449499). May also play a role as a coactivator for androgen- and, to a lesser extent, progesterone-dependent transcription (PubMed:19345326).

Cellular Location

Cytoplasm. Nucleus

Tissue Location


Widely expressed..


Anti-RNF14 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-RNF14 Rabbit Monoclonal Antibody - Images





Western blot analysis of RNF14 expression in Jurkat cell lysate.