

Anti-Meckelin Rabbit Monoclonal Antibody

Catalog # ABO16541

Specification

Anti-Meckelin Rabbit Monoclonal Antibody - Product Information

Application Primary Accession Host Isotype Reactivity Clonality Format **Description** WB, IHC, IF, ICC, FC <u>Q5HYA8</u> Rabbit IgG Rat, Human, Mouse Monoclonal Liquid

Anti-Meckelin Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-Meckelin Rabbit Monoclonal Antibody - Additional Information

Gene ID 91147

Other Names Meckelin, Meckel syndrome type 3 protein, Transmembrane protein 67, TMEM67, MKS3

Calculated MW 112 kDa KDa

Application Details WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
FC 1:50

Contents Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen A synthesized peptide derived from human Meckelin

Purification Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-Meckelin Rabbit Monoclonal Antibody - Protein Information

Name TMEM67



Synonyms MKS3

Function

Required for ciliary structure and function. Part of the tectonic-like complex which is required for tissue-specific ciliogenesis and may regulate ciliary membrane composition (By similarity). Involved in centrosome migration to the apical cell surface during early ciliogenesis. Involved in the regulation of cilia length and appropriate number through the control of centrosome duplication. Is a key regulator of stereociliary bundle orientation (By similarity). Required for epithelial cell branching morphology. Essential for endoplasmic reticulum-associated degradation (ERAD) of surfactant protein C (SFTPC). Involved in the negative regulation of canonical Wnt signaling, and activation of the non-canonical cascade stimulated by WNT5A (PubMed:26035863). In non-canonical Wnt signaling, it may act as ROR2 coreceptor (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein. Endoplasmic reticulum membrane; Multi-pass membrane protein. Cell projection, cilium Cytoplasm, cytoskeleton, cilium basal body. Note=Localizes at the transition zone, a region between the basal body and the ciliary axoneme (PubMed:22121117).

Tissue Location

Widely expressed in adult and fetal tissues. Expressed at higher level in spinal cord.

Anti-Meckelin Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Meckelin Rabbit Monoclonal Antibody - Images





All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.



Western blot analysis of Meckelin expression in (1) MCF7 cell lysate; (2) RAW 264.7 cell lysate; (3) C6 cell lysate.