

Anti-SOCS7 Rabbit Monoclonal Antibody

Catalog # ABO16561

Specification

Anti-SOCS7 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC
Primary Accession	<u>014512</u>
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	
Anti-SOCS7 Rabbit Monoclonal Antik	ody . Tested in WB, IHC applications. This antibody reacts with
Human.	

Anti-SOCS7 Rabbit Monoclonal Antibody - Additional Information

Gene ID 30837

Other Names Suppressor of cytokine signaling 7, SOCS-7, Nck, Ash and phospholipase C gamma-binding protein, Nck-associated protein 4, NAP-4, SOCS7, NAP4, SOCS6

Calculated MW 63 kDa KDa

Application Details WB 1:500-1:2000
HC 1:50-1:200

Contents Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen A synthesized peptide derived from human SOCS7

Purification Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-SOCS7 Rabbit Monoclonal Antibody - Protein Information

Name SOCS7 {ECO:0000303|PubMed:16127460, ECO:0000312|HGNC:HGNC:29846}



Function

Substrate-recognition component of a cullin-5-RING E3 ubiquitin-protein ligase complex (ECS complex, also named CRL5 complex), which mediates the ubiquitination and subsequent proteasomal degradation of target proteins, such as DAB1 and IRS1 (PubMed:16127460). Specifically recognizes and binds phosphorylated proteins via its SH2 domain, promoting their ubiquitination (By similarity). The ECS(SOCS7) complex acts as a key regulator of reelin signaling by mediating ubiquitination and degradation of phosphorylated DAB1 in the cortical plate of the developing cerebral cortex, thereby regulating neuron positioning during cortex development (By similarity). Functions in insulin signaling and glucose homeostasis through IRS1 ubiquitination and subsequent proteasomal degradation (PubMed:16127460" target="_blank">16127460" target="_blank">16127460" target="_blank">16127460" target="_blank">16127460" target="_blank">16127460" target="_blank">16127460). Also inhibits prolactin, growth hormone and leptin signaling by preventing STAT3 and STAT5 activation, sequestering them in the cytoplasm and reducing their binding to DNA (PubMed:15677474).

Cellular Location

Cytoplasm. Nucleus Cell membrane; Peripheral membrane protein; Cytoplasmic side. Note=Mostly cytoplasmic, but shuttles between the cytoplasm and the nucleus (PubMed:17803907). Rapidly relocalizes to the nucleus after UV irradiation (PubMed:17803907) Cytoplasmic location depends upon SEPT7 presence (PubMed:17803907)

Tissue Location

Expressed in brain and leukocytes (PubMed:9344857). Also in fetal lung fibroblasts and fetal brain (PubMed:9344857)

Anti-SOCS7 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-SOCS7 Rabbit Monoclonal Antibody - Images





Western blot analysis of SOCS7 expression in HeLa cell lysate.