

Anti-alpha Tubulin (acetyl K40) Rabbit Monoclonal Antibody
Catalog # ABO16708**Specification****Anti-alpha Tubulin (acetyl K40) Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IP, FC
Primary Accession	P68363/P68366
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-alpha Tubulin (acetyl K40) Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-alpha Tubulin (acetyl K40) Rabbit Monoclonal Antibody - Additional Information**Application Details**

WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
IP 1:50
FC 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human alpha Tubulin (acetyl K40)

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.


Anti-alpha Tubulin (acetyl K40) Rabbit Monoclonal Antibody - Protein Information**Anti-alpha Tubulin (acetyl K40) Rabbit Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-alpha Tubulin (acetyl K40) Rabbit Monoclonal Antibody - Images



Western blot analysis of alpha Tubulin (acetyl K40) expression in Mouse spleen cell lysate.



All lanes use the Antibody at 1:2W dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:2W dilution for 1 hour at room temperature.