

Anti-Phospho-alpha Synuclein (Y125) Rabbit Monoclonal Antibody

Catalog # ABO16794

Specification

Anti-Phospho-alpha Synuclein (Y125) Rabbit Monoclonal Antibody - Product Information

Application WB
Primary Accession P37840
Host Rabbit
Isotype Reactivity Human
Clonality Monoclonal
Format Liquid

Description

Anti-Phospho-alpha Synuclein (Y125) Rabbit Monoclonal Antibody . Tested in WB applications. This antibody reacts with Human.

Anti-Phospho-alpha Synuclein (Y125) Rabbit Monoclonal Antibody - Additional Information

Gene ID 6622

Other Names

Alpha-synuclein, Non-A beta component of AD amyloid, Non-A4 component of amyloid precursor, NACP, SNCA, NACP, PARK1

Application Details WB 1:500-1:2000

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunoaen

A synthesized peptide derived from human Phospho-alpha Synuclein (Y125)

Purification

Affinity-chromatography

Storage Store at -20°C for one year. For short term

storage and frequent use, store at 4°C for

up to one month. Avoid repeated

freeze-thaw cycles.

Anti-Phospho-alpha Synuclein (Y125) Rabbit Monoclonal Antibody - Protein Information

Name SNCA

Synonyms NACP, PARK1



Function

Neuronal protein that plays several roles in synaptic activity such as regulation of synaptic vesicle trafficking and subsequent neurotransmitter release (PubMed:<a href="http://www.uniprot.org/citations/20708282" target="http://www.uniprot.org/citations/20708282" target="http://www.uniprot.org/citations/207082

href="http://www.uniprot.org/citations/20798282" target="_blank">20798282, PubMed:26442590, PubMed:28288128, PubMed:30404828). Participates as a monomer in synaptic vesicle exocytosis by enhancing vesicle priming, fusion and dilation of exocytotic fusion pores (PubMed:28288128, PubMed:30404828). Mechanistically, acts by increasing local Ca(2+) release from microdomains which is essential for the enhancement of ATP-induced exocytosis (PubMed: 30404828). Also acts as a molecular chaperone in its multimeric membrane-bound state, assisting in the folding of synaptic fusion components called SNAREs (Soluble NSF Attachment Protein REceptors) at presynaptic plasma membrane in conjunction with cysteine string protein-alpha/DNAJC5 (PubMed:20798282). This chaperone activity is important to sustain normal SNARE-complex assembly during aging (PubMed:20798282). Also plays a role in the regulation of the dopamine neurotransmission by associating with the dopamine transporter (DAT1) and thereby modulating its activity (PubMed: 26442590).

Cellular Location

Cytoplasm. Membrane Nucleus Synapse. Secreted. Cell projection, axon {ECO:0000250|UniProtKB:O55042}. Note=Membrane-bound in dopaminergic neurons (PubMed:15282274). Expressed and colocalized with SEPTIN4 in dopaminergic axon terminals, especially at the varicosities (By similarity). {ECO:0000250|UniProtKB:O55042, ECO:0000269|PubMed:15282274}

Tissue Location

Highly expressed in presynaptic terminals in the central nervous system. Expressed principally in brain

Anti-Phospho-alpha Synuclein (Y125) Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Phospho-alpha Synuclein (Y125) Rabbit Monoclonal Antibody - Images