

**Anti-BTF Rabbit Monoclonal Antibody**  
**Catalog # ABO16810****Specification**

---

**Anti-BTF Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">Q9NYF8</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-BTF Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

**Anti-BTF Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 9774

**Other Names**

Bcl-2-associated transcription factor 1, Btf, BCLAF1 and THRAP3 family member 1, BCLAF1, BTF, KIAA0164

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human BTF

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-BTF Rabbit Monoclonal Antibody - Protein Information**

**Name** BCLAF1

**Synonyms** BTF, KIAA0164

**Function**

Death-promoting transcriptional repressor. May be involved in cyclin-D1/CCND1 mRNA stability through the SNARP complex which associates with both the 3'end of the CCND1 gene and its mRNA.

**Cellular Location**

Cytoplasm. Nucleus. Nucleus speckle. Nucleus, nucleoplasm


**Tissue Location**

Ubiquitous.

**Anti-BTF Rabbit Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-BTF Rabbit Monoclonal Antibody - Images**

All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.