

**p53 Antibody**  
**Rabbit Polyclonal Antibody**  
**Catalog # ABV10028****Specification**

---

**p53 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P04637</a>
Other Accession	<a href="#">EAW90140</a>
Reactivity	Human, Mouse, Rat, Rabbit, Dog
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	43653

**p53 Antibody - Additional Information****Gene ID** 7157

Application & Usage	Western blot analysis (0.5-4 µg/ml). It detects p53 in samples of human, mouse and rat origins. Other applications have not been determined.
---------------------	--

**Other Names**

TP53, TRP53, P53, LFS1, Tumor suppressor p53, Phosphoprotein p53

**Target/Specificity**

p53

**Antibody Form**

Liquid

**Appearance**

Colorless liquid

**Handling**

The antibody solution should be gently mixed before use.

**Reconstitution & Storage**

-20 °C

**Background Descriptions****Precautions**

p53 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**p53 Antibody - Protein Information**

**Name** TP53**Synonyms** P53**Function**

Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type (PubMed:<a

href="http://www.uniprot.org/citations/11025664" target="\_blank">11025664</a>, PubMed:<a href="http://www.uniprot.org/citations/12524540" target="\_blank">12524540</a>, PubMed:<a href="http://www.uniprot.org/citations/12810724" target="\_blank">12810724</a>, PubMed:<a href="http://www.uniprot.org/citations/15186775" target="\_blank">15186775</a>, PubMed:<a href="http://www.uniprot.org/citations/15340061" target="\_blank">15340061</a>, PubMed:<a href="http://www.uniprot.org/citations/17317671" target="\_blank">17317671</a>, PubMed:<a href="http://www.uniprot.org/citations/17349958" target="\_blank">17349958</a>, PubMed:<a href="http://www.uniprot.org/citations/19556538" target="\_blank">19556538</a>, PubMed:<a href="http://www.uniprot.org/citations/20673990" target="\_blank">20673990</a>, PubMed:<a href="http://www.uniprot.org/citations/20959462" target="\_blank">20959462</a>, PubMed:<a href="http://www.uniprot.org/citations/22726440" target="\_blank">22726440</a>, PubMed:<a href="http://www.uniprot.org/citations/24051492" target="\_blank">24051492</a>, PubMed:<a href="http://www.uniprot.org/citations/9840937" target="\_blank">9840937</a>, PubMed:<a href="http://www.uniprot.org/citations/24652652" target="\_blank">24652652</a>). Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process (PubMed:<a href="http://www.uniprot.org/citations/11025664" target="\_blank">11025664</a>, PubMed:<a href="http://www.uniprot.org/citations/12524540" target="\_blank">12524540</a>, PubMed:<a href="http://www.uniprot.org/citations/12810724" target="\_blank">12810724</a>, PubMed:<a href="http://www.uniprot.org/citations/15186775" target="\_blank">15186775</a>, PubMed:<a href="http://www.uniprot.org/citations/15340061" target="\_blank">15340061</a>, PubMed:<a href="http://www.uniprot.org/citations/17317671" target="\_blank">17317671</a>, PubMed:<a href="http://www.uniprot.org/citations/17349958" target="\_blank">17349958</a>, PubMed:<a href="http://www.uniprot.org/citations/19556538" target="\_blank">19556538</a>, PubMed:<a href="http://www.uniprot.org/citations/20673990" target="\_blank">20673990</a>, PubMed:<a href="http://www.uniprot.org/citations/20959462" target="\_blank">20959462</a>, PubMed:<a href="http://www.uniprot.org/citations/22726440" target="\_blank">22726440</a>, PubMed:<a href="http://www.uniprot.org/citations/24051492" target="\_blank">24051492</a>, PubMed:<a href="http://www.uniprot.org/citations/9840937" target="\_blank">9840937</a>, PubMed:<a href="http://www.uniprot.org/citations/24652652" target="\_blank">24652652</a>). One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. Its pro-apoptotic activity is activated via its interaction with PPP1R13B/ASPP1 or TP53BP2/ASPP2 (PubMed:<a href="http://www.uniprot.org/citations/12524540" target="\_blank">12524540</a>). However, this activity is inhibited when the interaction with PPP1R13B/ASPP1 or TP53BP2/ASPP2 is displaced by PPP1R13L/iASPP (PubMed:<a href="http://www.uniprot.org/citations/12524540" target="\_blank">12524540</a>). In cooperation with mitochondrial PPIF is involved in activating oxidative stress-induced necrosis; the function is largely independent of transcription. Induces the transcription of long intergenic non-coding RNA p21 (lincRNA-p21) and lincRNA-Mkln1. LincRNA-p21 participates in TP53-dependent transcriptional repression leading to apoptosis and seems to have an effect on cell-cycle regulation. Implicated in Notch signaling cross-over. Prevents CDK7 kinase activity when associated to CAK complex in response to DNA damage, thus stopping cell cycle progression. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediated apoptosis. Regulates the circadian clock by repressing CLOCK-BMAL1-mediated transcriptional activation of PER2 (PubMed:<a href="http://www.uniprot.org/citations/24051492" target="\_blank">24051492</a>).

**Cellular Location**

Cytoplasm. Nucleus. Nucleus, PML body Endoplasmic reticulum. Mitochondrion matrix. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Note=Recruited into PML bodies together with CHEK2 (PubMed:12810724) Translocates to mitochondria upon oxidative stress (PubMed:22726440) Translocates to mitochondria in response to mitomycin C treatment (PubMed:27323408). Competitive inhibition of TP53 interaction with HSPA9/MOT-2 by UBXN2A results in increased protein abundance and subsequent translocation of TP53 to the nucleus (PubMed:24625977) [Isoform 2]: Nucleus. Cytoplasm. Note=Localized mainly in the nucleus with minor staining in the cytoplasm [Isoform 4]: Nucleus. Cytoplasm. Note=Predominantly nuclear but translocates to the cytoplasm following cell stress [Isoform 8]: Nucleus. Cytoplasm. Note=Localized in both nucleus and cytoplasm in most cells. In some cells, forms foci in the nucleus that are different from nucleoli

### Tissue Location


Ubiquitous. Isoforms are expressed in a wide range of normal tissues but in a tissue-dependent manner. Isoform 2 is expressed in most normal tissues but is not detected in brain, lung, prostate, muscle, fetal brain, spinal cord and fetal liver. Isoform 3 is expressed in most normal tissues but is not detected in lung, spleen, testis, fetal brain, spinal cord and fetal liver. Isoform 7 is expressed in most normal tissues but is not detected in prostate, uterus, skeletal muscle and breast. Isoform 8 is detected only in colon, bone marrow, testis, fetal brain and intestine. Isoform 9 is expressed in most normal tissues but is not detected in brain, heart, lung, fetal liver, salivary gland, breast or intestine

### p53 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### p53 Antibody - Images



Western blot analysis of p53 in 3T3 (Lane 1) and Jurkat (Lane 2, 3) cell lysates.

### p53 Antibody - Background

p53, a 53 kDa protein, binds to a DNA consensus sequence, the p53 response element, and regulates normal cell cycle events by activating transcription of genes involved either in progression through the cell cycle, or causing arrest in G1 when the genome is damaged. In most transformed and tumor cells the concentration of p53 is increased 5-1000 fold over the concentration in normal cells, principally due to the increased half-life (4 hrs) compared to that of wild-type (20 min). p53 localizes in the nucleus, but is detectable at the plasma membrane during mitosis and certain mutations also modulate cytoplasmic/nuclear distribution. p53 downregulates Bcl-2 expression and upregulates Bax expression, but may not always be necessary for apoptosis.