

CREB Antibody

Rabbit Polyclonal Antibody Catalog # ABV10263

Specification

CREB Antibody - Product Information

Application WB
Primary Accession P15337

Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 35081

CREB Antibody - Additional Information

Gene ID 81646

Application & Usage Western blotting (0.5-4 μg/ml). However,

the optimal concentrations should be determined individually. The antibody recognizes 43 kDa CREB from samples of

human, mouse and rat origins.

Other Names

CREB-1, cAMP-response Element-binding Protein-1, CREB1, CREB, MGC9284

Target/Specificity

CREB

Antibody Form

Liquid

Appearance

Colorless liquid

Formulation

 $100~\mu g$ (0.5 mg/ml) purified rabbit polyclonal antibody in phosphate-buffered saline (PBS) containing 30% glycerol, 0.5% BSA and 0.01% thimerosal.

Handling

The antibody solution should be gently mixed before use.

Reconstitution & Storage

-20 °C

Background Descriptions

Precautions

CREB Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



CREB Antibody - Protein Information

Name Creb1

Synonyms Creb-1

Function

Phosphorylation-dependent transcription factor that stimulates transcription upon binding to the DNA cAMP response element (CRE), a sequence present in many viral and cellular promoters (By similarity). Transcription activation is enhanced by the TORC coactivators which act independently of Ser-119 phosphorylation (By similarity). Involved in different cellular processes including the synchronization of circadian rhythmicity and the differentiation of adipose cells (By similarity). Regulates the expression of apoptotic and inflammatory response factors in cardiomyocytes in response to ERFE-mediated activation of AKT signaling (PubMed:https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056">https://www.uniprot.org/citations/30566056

Cellular Location Nucleus.

CREB Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CREB Antibody - Images

CREB Antibody - Background

Transcription factor CREB binds the cAMP response element (CRE) and activates transcription in response to a variety of extracellular signals including neurotransmitters, hormones, membrane depolarization, and growth or neurotrophic factors. Protein kinase A and the calmodulin-dependent protein kinase CaMKII stimulate CREB phosphorylation at Ser133, a key regulatory site controlling transcriptional activity. Phosphorylation at Ser133 is also mediated by p44/42 MAP kinase, p90RSK, p38 MAP kinase and MSK1. CREB appears to play an important role in learning and memory in both flies and mice.