

Cox-2 Antibody
Rabbit Polyclonal Antibody
Catalog # ABV10265

Specification

Cox-2 Antibody - Product Information

Application	WB
Primary Accession	P35354
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	68996

Cox-2 Antibody - Additional Information

Gene ID 5743

Application & Usage

Western blotting (0.5-4 µg/ml). However, the optimal concentrations should be determined individually. The antibody recognizes 66 kDa human Cox-2. It does not cross-react with Cox-1. Jurkat cell lysate can be used as a positive control.

Other Names

Cyclooxygenase 2, PTGS2, PHS-2, COX-2, hCox-2 , PGHS-2, PGG/HS, COX2, EC 1.14.99.1

Target/Specificity

Cox-2

Antibody Form

Liquid

Appearance

Colorless liquid

Formulation

100 µg (0.5 mg/ml) affinity purified rabbit polyclonal antibody in phosphate-buffered saline (PBS) containing 30% glycerol, 0.5% BSA, and 0.01% thimerosal.

Handling

The antibody solution should be gently mixed before use.

Reconstitution & Storage

-20 °C

Background Descriptions

Precautions

Cox-2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Cox-2 Antibody - Protein Information

Name PTGS2 ([HGNC:9605](#))

Function

Dual cyclooxygenase and peroxidase in the biosynthesis pathway of prostanoids, a class of C20 oxylipins mainly derived from arachidonate ((5Z,8Z,11Z,14Z)-eicosatetraenoate, AA, C20:4(n-6)), with a particular role in the inflammatory response (PubMed:<a href="<http://www.uniprot.org/citations/7947975>">7947975, PubMed:<a href="<http://www.uniprot.org/citations/7592599>">7592599, PubMed:<a href="<http://www.uniprot.org/citations/9261177>">9261177, PubMed:<a href="<http://www.uniprot.org/citations/16373578>">16373578, PubMed:<a href="<http://www.uniprot.org/citations/22942274>">22942274, PubMed:<a href="<http://www.uniprot.org/citations/26859324>">26859324, PubMed:<a href="<http://www.uniprot.org/citations/27226593>">27226593, PubMed:<a href="<http://www.uniprot.org/citations/11939906>">11939906, PubMed:<a href="<http://www.uniprot.org/citations/19540099>">19540099). The cyclooxygenase activity oxygenates AA to the hydroperoxy endoperoxide prostaglandin G2 (PGG2), and the peroxidase activity reduces PGG2 to the hydroxy endoperoxide prostaglandin H2 (PGH2), the precursor of all 2-series prostaglandins and thromboxanes (PubMed:<a href="<http://www.uniprot.org/citations/7947975>">7947975, PubMed:<a href="<http://www.uniprot.org/citations/7592599>">7592599, PubMed:<a href="<http://www.uniprot.org/citations/9261177>">9261177, PubMed:<a href="<http://www.uniprot.org/citations/16373578>">16373578, PubMed:<a href="<http://www.uniprot.org/citations/22942274>">22942274, PubMed:<a href="<http://www.uniprot.org/citations/26859324>">26859324, PubMed:<a href="<http://www.uniprot.org/citations/27226593>">27226593). This complex transformation is initiated by abstraction of hydrogen at carbon 13 (with S-stereochemistry), followed by insertion of molecular O₂ to form the endoperoxide bridge between carbon 9 and 11 that defines prostanoids. The insertion of a second molecule of O₂ (bis-oxygenase activity) yields a hydroperoxy group in PGG2 that is then reduced to PGH2 by two electrons (PubMed:<a href="<http://www.uniprot.org/citations/7947975>">7947975, PubMed:<a href="<http://www.uniprot.org/citations/7592599>">7592599, PubMed:<a href="<http://www.uniprot.org/citations/9261177>">9261177, PubMed:<a href="<http://www.uniprot.org/citations/16373578>">16373578, PubMed:<a href="<http://www.uniprot.org/citations/22942274>">22942274, PubMed:<a href="<http://www.uniprot.org/citations/26859324>">26859324, PubMed:<a href="<http://www.uniprot.org/citations/27226593>">27226593). Similarly catalyzes successive cyclooxygenation and peroxidation of dihomo-gamma-linoleate (DGLA, C20:3(n-6)) and eicosapentaenoate (EPA, C20:5(n-3)) to corresponding PGH1 and PGH3, the precursors of 1- and 3- series prostaglandins (PubMed:<a href="<http://www.uniprot.org/citations/11939906>">11939906, PubMed:<a href="<http://www.uniprot.org/citations/19540099>">19540099). In an alternative pathway of prostanoid biosynthesis, converts 2-arachidonoyl lysophospholipids to prostanoid lysophospholipids, which are then hydrolyzed by intracellular phospholipases to release free prostanoids (PubMed:27642067). Metabolizes 2-arachidonoyl glycerol yielding the glyceryl ester of PGH2, a process that can contribute to pain response (PubMed:22942274). Generates lipid mediators from n-3 and n-6 polyunsaturated fatty acids (PUFAs) via a lipoxygenase-type mechanism. Oxygenates PUFAs to hydroperoxy compounds and then reduces them to corresponding alcohols (PubMed:11034610, PubMed:11192938).

target="_blank">>11192938, PubMed:9048568, PubMed:9261177). Plays a role in the generation of resolution phase interaction products (resolvins) during both sterile and infectious inflammation (PubMed:12391014). Metabolizes docosahexaenoate (DHA, C22:6(n-3)) to 17R-HDHA, a precursor of the D- series resolvins (RvDs) (PubMed:12391014). As a component of the biosynthetic pathway of E-series resolvins (RvEs), converts eicosapentaenoate (EPA, C20:5(n-3)) primarily to 18S-HEPE that is further metabolized by ALOX5 and LTA4H to generate 18S-RvE1 and 18S- RvE2 (PubMed:21206090). In vascular endothelial cells, converts docosapentaenoate (DPA, C22:5(n-3)) to 13R-HDPA, a precursor for 13-series resolvins (RvTs) shown to activate macrophage phagocytosis during bacterial infection (PubMed:26236990). In activated leukocytes, contributes to oxygenation of hydroxyeicosatetraenoates (HETE) to diHETES (5,15-diHETE and 5,11-diHETE) (PubMed:22068350, PubMed:26282205). Can also use linoleate (LA, (9Z,12Z)- octadecadienoate, C18:2(n-6)) as substrate and produce hydroxyoctadecadienoates (HODEs) in a regio- and stereospecific manner, being (9R)-HODE ((9R)-hydroxy-(10E,12Z)-octadecadienoate) and (13S)- HODE ((13S)-hydroxy-(9Z,11E)-octadecadienoate) its major products (By similarity). During neuroinflammation, plays a role in neuronal secretion of specialized preresolving mediators (SPMs) 15R-lipoxin A4 that regulates phagocytic microglia (By similarity).

Cellular Location

Microsome membrane; Peripheral membrane protein. Endoplasmic reticulum membrane; Peripheral membrane protein. Nucleus inner membrane; Peripheral membrane protein. Nucleus outer membrane; Peripheral membrane protein. Note=Detected on the luminal side of the endoplasmic reticulum and nuclear envelope

Cox-2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Cox-2 Antibody - Images

Cox-2 Antibody - Background

COX proteins are membrane-associated heme proteins that have cyclooxygenase and peroxidase activities. These enzymes are targets of NSAID (Nonsteroidal anti-inflammatory dr µgs) such as aspirin. Prostaglandins (PGs) formed by the enzymatic activity of COX-1 are primarily involved in the regulation of homeostatic functions thro µghout the body, whereas PGs formed by COX-2 primarily mediate pain, fever, and inflammation. COX-1 is constitutively expressed, with particularly high expression in gastrointestinal tissues. COX-2 is induced by cytokines and mitogens and is likely to play a role in inflammatory diseases such as rheumatoid arthritis.