

**LIMK1 Antibody**  
**Rabbit Polyclonal Antibody**  
**Catalog # ABV10315**

**Specification**

---

**LIMK1 Antibody - Product Information**

Application	WB, IP
Primary Accession	<a href="#">P53667</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	72585

**LIMK1 Antibody - Additional Information**

**Gene ID** 3984

**Application & Usage**

Western blotting (1-4 µg/ml). However, the optimal conditions should be determined individually. The antibody detects 70 kDa LIMK1 protein. It does not cross-react with LIMK2.

**Other Names**

LIM kinases, LIMK-1

**Target/Specificity**

LIMK1

**Antibody Form**

Liquid

**Appearance**

Colorless liquid

**Formulation**

100 µg (0.5 mg/ml) antigen affinity purified rabbit polyclonal antibody in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA, 0.02% thimerosal.

**Handling**

The antibody solution should be gently mixed before use.

**Reconstitution & Storage**

-20 °C

**Background Descriptions**

**Precautions**

LIMK1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## LIMK1 Antibody - Protein Information

**Name** LIMK1

**Synonyms** LIMK

### Function

Serine/threonine-protein kinase that plays an essential role in the regulation of actin filament dynamics. Acts downstream of several Rho family GTPase signal transduction pathways (PubMed:<a href="http://www.uniprot.org/citations/10436159" target="\_blank">10436159</a>, PubMed:<a href="http://www.uniprot.org/citations/11832213" target="\_blank">11832213</a>, PubMed:<a href="http://www.uniprot.org/citations/12807904" target="\_blank">12807904</a>, PubMed:<a href="http://www.uniprot.org/citations/15660133" target="\_blank">15660133</a>, PubMed:<a href="http://www.uniprot.org/citations/16230460" target="\_blank">16230460</a>, PubMed:<a href="http://www.uniprot.org/citations/18028908" target="\_blank">18028908</a>, PubMed:<a href="http://www.uniprot.org/citations/22328514" target="\_blank">22328514</a>, PubMed:<a href="http://www.uniprot.org/citations/23633677" target="\_blank">23633677</a>). Activated by upstream kinases including ROCK1, PAK1 and PAK4, which phosphorylate LIMK1 on a threonine residue located in its activation loop (PubMed:<a href="http://www.uniprot.org/citations/10436159" target="\_blank">10436159</a>). LIMK1 subsequently phosphorylates and inactivates the actin binding/depolymerizing factors cofilin-1/CFL1, cofilin-2/CFL2 and destin/DSTN, thereby preventing the cleavage of filamentous actin (F-actin), and stabilizing the actin cytoskeleton (PubMed:<a href="http://www.uniprot.org/citations/11832213" target="\_blank">11832213</a>, PubMed:<a href="http://www.uniprot.org/citations/15660133" target="\_blank">15660133</a>, PubMed:<a href="http://www.uniprot.org/citations/16230460" target="\_blank">16230460</a>, PubMed:<a href="http://www.uniprot.org/citations/23633677" target="\_blank">23633677</a>). In this way LIMK1 regulates several actin-dependent biological processes including cell motility, cell cycle progression, and differentiation (PubMed:<a href="http://www.uniprot.org/citations/11832213" target="\_blank">11832213</a>, PubMed:<a href="http://www.uniprot.org/citations/15660133" target="\_blank">15660133</a>, PubMed:<a href="http://www.uniprot.org/citations/16230460" target="\_blank">16230460</a>, PubMed:<a href="http://www.uniprot.org/citations/23633677" target="\_blank">23633677</a>). Phosphorylates TPPP on serine residues, thereby promoting microtubule disassembly (PubMed:<a href="http://www.uniprot.org/citations/18028908" target="\_blank">18028908</a>). Stimulates axonal outgrowth and may be involved in brain development (PubMed:<a href="http://www.uniprot.org/citations/18028908" target="\_blank">18028908</a>).

### Cellular Location

Cytoplasm. Nucleus. Cytoplasm, cytoskeleton. Cell projection, lamellipodium {ECO:0000250|UniProtKB:P53668} Note=Predominantly found in the cytoplasm. Localizes in the lamellipodium in a CDC42BPA, CDC42BPB and FAM89B/LRAP25-dependent manner. {ECO:0000250|UniProtKB:P53668}

### Tissue Location

Highest expression in both adult and fetal nervous system. Detected ubiquitously throughout the different regions of adult brain, with highest levels in the cerebral cortex. Expressed to a lesser extent in heart and skeletal muscle

## LIMK1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### LIMK1 Antibody - Images

### LIMK1 Antibody - Background

LIM kinases (LIMK1 and LIMK2) are serine/threonine kinases that have two zinc finger motifs, known as LIM motifs in their N-terminal regulatory domain. LIM kinases are involved in actin cytoskeleton regulation through Rho-family GTPases and downstream kinases PAKs and ROCK. PAK1 and ROCK phosphorylate LIMK1 and LIMK2, which increases the activity of the kinases. Activated LIM kinases inhibit the actin depolymerization activity of cofilin by phosphorylation at the N-terminus of Cofilin.