

G3BP Antibody
Rabbit Polyclonal Antibody
Catalog # ABV10464

Specification

G3BP Antibody - Product Information

Application	WB
Primary Accession	Q13283
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	52164

G3BP Antibody - Additional Information

Gene ID 10146

Positive Control

Western blot: Jurkat, 3T3, rat kidney cell lysate

Application & Usage

Western blot: 1:200

Other Names

ATP-dependent DNA helicase VIII

Target/Specificity

G3BP

Antibody Form

Liquid

Appearance

Colorless liquid

Formulation

100 µg (0.5 mg/ml) of antibody in PBS, 0.01 % BSA, 0.01 % thimerosal, and 50 % glycerol, pH 7.2

Handling

The antibody solution should be gently mixed before use.

Reconstitution & Storage

-20 °C

Background Descriptions

Precautions

G3BP Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

G3BP Antibody - Protein Information

Name G3BP1 {ECO:0000303|PubMed:23279204, ECO:0000312|HGNC:HGNC:30292}

Function

Protein involved in various processes, such as stress granule formation and innate immunity (PubMed:12642610, PubMed:20180778, PubMed:23279204, PubMed:30510222, PubMed:30804210). Plays an essential role in stress granule formation (PubMed:12642610, PubMed:20180778, PubMed:23279204, PubMed:32302570, PubMed:32302571, PubMed:32302572, PubMed:36183834, PubMed:36279435, PubMed:34739333, PubMed:36692217, PubMed:37379838). Stress granules are membraneless compartments that store mRNAs and proteins, such as stalled translation pre-initiation complexes, in response to stress (PubMed:12642610, PubMed:20180778, PubMed:23279204, PubMed:27022092, PubMed:32302570, PubMed:32302571, PubMed:32302572, PubMed:36279435, PubMed:37379838). Promotes formation of stress granules phase-separated membraneless compartment by undergoing liquid-liquid phase separation (LLPS) upon unfolded RNA- binding: functions as a molecular switch that triggers RNA-dependent LLPS in response to a rise in intracellular free RNA concentrations (PubMed:32302570, PubMed:32302571, PubMed:32302572, PubMed:34739333, PubMed:36692217, PubMed:36279435). Also acts as an ATP- and magnesium- dependent helicase: unwinds DNA/DNA, RNA/DNA, and RNA/RNA substrates with comparable efficiency (PubMed:9889278). Acts unidirectionally by moving in the 5' to 3' direction along the bound single-stranded DNA (PubMed:9889278). Unwinds preferentially partial DNA and RNA duplexes having a 17 bp annealed portion and either a hanging 3' tail or hanging tails at both 5'- and 3'-ends (PubMed:9889278). Plays an essential role in innate immunity by promoting CGAS and RIGI activity (PubMed:30510222, PubMed:30804210). Participates in the DNA-triggered cGAS/STING pathway by promoting the DNA binding and activation of CGAS (PubMed:30510222). Triggers the condensation of cGAS, a process probably linked to the formation of membrane-less organelles (PubMed:34779554)

target="_blank">34779554). Enhances also RIGI-induced type I interferon production probably by helping RIGI at sensing pathogenic RNA (PubMed:30804210). May also act as a phosphorylation-dependent sequence-specific endoribonuclease in vitro: Cleaves exclusively between cytosine and adenine and cleaves MYC mRNA preferentially at the 3'-UTR (PubMed:11604510).

Cellular Location

Cytoplasm, cytosol. Perikaryon {ECO:0000250|UniProtKB:P97855}. Cytoplasm, Stress granule. Nucleus Note=Cytoplasmic in proliferating cells (PubMed:11604510). Cytosolic and partially nuclear in resting cells (PubMed:11604510). Recruited to stress granules in response to arsenite treatment (PubMed:12642610, PubMed:20180778). The unphosphorylated form is recruited to stress granules (PubMed:12642610). HRAS signaling contributes to this process by regulating G3BP dephosphorylation (PubMed:12642610)

Tissue Location

Ubiquitous..

G3BP Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

G3BP Antibody - Images

G3BP Antibody - Background

G3BP1 (GTPase activating protein (SH3 domain) binding protein 1), also known as G3BP or HDH-VIII, is a ubiquitously expressed protein that localizes to the cytoplasm in proliferating cells and to the nucleus in non-proliferating cells. One of several DNA-unwinding enzymes, G3BP1 functions as a sequence-specific, phosphorylation-dependent helicase that unwinds partial RNA and DNA duplexes containing hanging 3'- or 5'- ends. G3BP1 uses magnesium as a cofactor and, in addition to its helicase activity, acts as an endoribonuclease that cleaves mRNA between adenine and cytosine residues at the 3'-UTR. An element of the Ras signaling pathway, G3BP1 binds to the SH3 domain of Ras GTPase-activating protein (Ras GAP) in proliferating cells, thereby regulating Ras signaling events in developing tissues. Due to its involvement in both DNA replication and signaling pathways within the cell, G3BP1 expression is implicated in the pathogenesis of several cancers, including esophageal squamous carcinoma.