

**MCM2 Antibody**  
**Rabbit Polyclonal Antibody**  
**Catalog # ABV10577****Specification**

---

**MCM2 Antibody - Product Information**

|                   |                             |
|-------------------|-----------------------------|
| Application       | WB, IP                      |
| Primary Accession | <a href="#">P49736</a>      |
| Other Accession   | <a href="#">NP_004511.2</a> |
| Reactivity        | Human, Mouse                |
| Host              | Rabbit                      |
| Clonality         | Polyclonal                  |
| Isotype           | Rabbit IgG                  |
| Calculated MW     | 101896                      |

**MCM2 Antibody - Additional Information****Gene ID** 4171

|                     |  |
|---------------------|--|
| Application & Usage | <b>Western blotting (1:500 - 1:2500) and Immunoprecipitation. HeLa cell extract can be used as a positive control. However, the optimal concentrations should be determined individually. The antibody recognizes the MCM2 of human origin. Reactivity to mouse is expected due to sequence homology. Reactivity to other species has not been tested.</b> |
|---------------------|--|

**Other Names**

Mitotin; BM28, Nuclear protein BM28; cdc19; CCNL1, cyclin like-1; CDCL1, CDC like-1; D3S3194; KIAA0030

**Target/Specificity**

MCM2

**Antibody Form**

Liquid

**Appearance**

Colorless liquid

**Formulation**

100 µl affinity purified rabbit polyclonal antibody in phosphate-buffered saline (PBS) containing 30% glycerol, 1% BSA and 0.02% thimerosal.

**Handling**

The antibody solution should be gently mixed before use.

**Reconstitution & Storage**

-20 °C

## Background Descriptions

### Precautions

MCM2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## MCM2 Antibody - Protein Information

**Name** MCM2 ([HGNC:6944](#))

### Function

Acts as a component of the MCM2-7 complex (MCM complex) which is the replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. Core component of CDC45-MCM-GINS (CMG) helicase, the molecular machine that unwinds template DNA during replication, and around which the replisome is built (PubMed:<a href="http://www.uniprot.org/citations/32453425" target="\_blank">32453425</a>, PubMed:<a href="http://www.uniprot.org/citations/34694004" target="\_blank">34694004</a>, PubMed:<a href="http://www.uniprot.org/citations/34700328" target="\_blank">34700328</a>, PubMed:<a href="http://www.uniprot.org/citations/35585232" target="\_blank">35585232</a>). The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity (PubMed:<a href="http://www.uniprot.org/citations/32453425" target="\_blank">32453425</a>). Required for the entry in S phase and for cell division (PubMed:<a href="http://www.uniprot.org/citations/8175912" target="\_blank">8175912</a>). Plays a role in terminally differentiated hair cells development of the cochlea and induces cells apoptosis (PubMed:<a href="http://www.uniprot.org/citations/26196677" target="\_blank">26196677</a>).

### Cellular Location

Nucleus. Chromosome. Note=Associated with chromatin before the formation of nuclei and detaches from it as DNA replication progresses. {ECO:0000250|UniProtKB:P55861}

## MCM2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## MCM2 Antibody - Images

## MCM2 Antibody - Background

The mini-chromosome maintenance (MCM) family of proteins, including MCM2, MCM3, MCM4 (Cdc21), MCM5 (Cdc46), MCM6 (Mis5) and MCM7 (Cdc47), are regulators of DNA replication that act to ensure replication occurs only once in the cell cycle. Expression of MCM proteins increases during cell growth, peaking at G1 to S phase. The MCM proteins each contain an ATP-binding motif, which

is predicted to mediate ATP-dependent opening of double-stranded DNA. MCM proteins are regulated by E2F transcription factors, which induce MCM expression, and by protein kinases, which interact with MCM proteins to maintain the postreplicative state of the cell. MCM2/MCM4 complexes function as substrates for Cdc2/cyclin B in vitro. Cleavage of MCM3, which can be prevented by caspase inhibitors, results in the inactivation during apoptosis of the MCM complex, which is composed of, at least, MCM2-6. A complex composed of MCM4, MCM6 and MCM7 has been shown to be involved in DNA helicase activity, and MCM5 is involved in IFN- $\gamma$ -induced Stat1 $\alpha$  transcription activation.