

Matrilin-2 Antibody

Rabbit Polyclonal Antibody Catalog # ABV10651

Specification

Matrilin-2 Antibody - Product Information

Application Primary Accession Reactivity Host Clonality

Isotype Calculated MW WB

000339

Human, Mouse, Rat

Rabbit Polyclonal Rabbit IgG 106837

Matrilin-2 Antibody - Additional Information

Gene ID 4147

Application & Usage

Western blotting (0.5-4 μ g/ml). However, the optimal concentrations should be determined individually. The antibody recognizes 120-150 kDa Matrilin-2 from samples of human, mouse and rat origins. An additional ~55 kDa band and a ~38 kDa band can also be detected in rat kidney and Jurkat cells, individually. Other species have not been determined.

Other Names MATN2

Target/Specificity Matrilin-2

Antibody Form Liquid

AppearanceColorless liquid

Formulation

 $100 \mu g$ (0.5 mg/ml) affinity purified rabbit anti-Matrilin-2 polyclonal antibody in phosphate-buffered saline (PBS) containing 30% glycerol, 0.5% BSA, and 0.01% thimerosal.

Handling

The antibody solution should be gently mixed before use.

Reconstitution & Storage -20 °C

Background Descriptions



Tel: 858.875.1900 Fax: 858.875.1999

Precautions

Matrilin-2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Matrilin-2 Antibody - Protein Information

Name MATN2

Function

Involved in matrix assembly.

Cellular Location


Secreted.

Matrilin-2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Matrilin-2 Antibody - Images



Western blot analysis of Matrilin-2 in Rat kidney tissue lysate (Lane 1), 3T3 cell lysate (Lane 2) and Jurkat cell lysate (Lane 3).

Matrilin-2 Antibody - Background

Matrilin-2 is part of the Matrilin family of secreted extracelluar matrix proteins that consist of Matrilin-1 to Matrilin-4. Matrilin-2 encodes a member of the von Willebrand factor A domain protein





family. It is involved in matrix assembly and development of filamentous networks in the extracellular matrices of various tissues.