

S-Tag Antibody
Rabbit Polyclonal Antibody
Catalog # ABV10758**Specification**

S-Tag Antibody - Product InformationApplication
Host
Clonality
Isotype**WB**
Rabbit
Polyclonal
Rabbit IgG**S-Tag Antibody - Additional Information**

Application & Usage

The antibody can be used for Western blot analysis (0.5-4 µg/ml). However, the optimal conditions should be determined individually.**Other Names**

S-tag, anti-S-tag, S-tag antibody

Target/Specificity

S-Tag

Antibody Form

Liquid

Appearance

Colorless liquid

Formulation

100 µg (0.5 mg/ml) affinity purified rabbit anti-S-tag polyclonal antibody in phosphate buffered saline (PBS), pH 7.2, containing 30% glycerol, 0.5% BSA, 5mM EDTA and 0.01% thimerosal.

Handling

The antibody solution should be gently mixed before use.

Reconstitution & Storage

-20 °C

Background Descriptions**Precautions**

S-Tag Antibody is for research use only and not for use in diagnostic or therapeutic procedures.


S-Tag Antibody - Protein Information

S-Tag Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

S-Tag Antibody - Images



Western blot analysis using E.coli lysate expressing S-Tag .

S-Tag Antibody - Background

S-tag is an epitope tag composed of a 15 residue peptide derived from pancreatic ribonuclease A