

SLC Antibody
Rabbit Polyclonal Antibody
Catalog # ABV10987**Specification**

SLC Antibody - Product Information

Application	WB
Primary Accession	P84443
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG

SLC Antibody - Additional Information

Application & Usage	Western blot analysis (0.5-1 µg/ml). However, the optimal conditions should be determined individually. Cross-reactivity to SLC of other species has not been determined.
---------------------	--

Other Names

Secondary lymphoid-tissue chemokine, TCA4, 6Ckine, Exodus-2, Exodus 2, Exodus2, C-C motif chemokine 21, Small-inducible cytokine A21

Target/Specificity

SLC

Antibody Form

Liquid

Appearance

Colorless liquid

Formulation

100 µg (0.5 mg/ml) Protein A purified rabbit anti-rat SLC polyclonal antibody in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA, 0.02% sodium azide.

Handling

The antibody solution should be gently mixed before use.

Reconstitution & Storage

-20 °C

Background Descriptions**Precautions**

SLC Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

SLC Antibody - Protein Information

SLC Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SLC Antibody - Images

SLC Antibody - Background

SLC (secondary lymphoid-tissue chemokine, also called 6CKine, TCA4, Exodus-2) is a 111-amino acid (human; 110 a.a. for mouse) CC chemokine with an N-terminal sequence of 23 residues as a signal sequence. It was originally cloned in 1997. Recombinant SLC attracts T-, B- cells, mature dendritic cells, but did not induce any cell migration in neutrophils or monocytes. SLC is a functional ligand for CCR7.