

**ApoE3 Antibody**  
**Rabbit Polyclonal Antibody**  
**Catalog # ABV11050****Specification**

---

**ApoE3 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P02649</a>
Other Accession	<a href="#">NP_000032</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	36154

**ApoE3 Antibody - Additional Information****Gene ID** 348**Application & Usage**

**Western blotting (0.5-4 µg/ml). However, the optimal concentrations should be determined individually. Recombinant ApoA1 can be used as a positive control. The antibody recognizes ~34 kDa of human ApoE3. Reactivity to other species has not been tested.**

**Other Names**

Apolipoprotein E3, APOE 3, APOE3, APOE-3

**Target/Specificity**

ApoE3

**Antibody Form**

Liquid

**Appearance**

Colorless liquid

**Formulation**

100 µg (0.5 mg/ml) affinity purified rabbit polyclonal antibody in phosphate-buffered saline (PBS), pH 7.2, containing 30% glycerol, 0.5% BSA, and 0.01% thimerosal.

**Handling**

The antibody solution should be gently mixed before use.

**Reconstitution & Storage**

-20 °C

## Background Descriptions

### Precautions

ApoE3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## ApoE3 Antibody - Protein Information

**Name** APOE ([HGNC:613](#))

### Function

APOE is an apolipoprotein, a protein associating with lipid particles, that mainly functions in lipoprotein-mediated lipid transport between organs via the plasma and interstitial fluids (PubMed:<a href="http://www.uniprot.org/citations/6860692" target="\_blank">6860692</a>, PubMed:<a href="http://www.uniprot.org/citations/1911868" target="\_blank">1911868</a>, PubMed:<a href="http://www.uniprot.org/citations/14754908" target="\_blank">14754908</a>). APOE is a core component of plasma lipoproteins and is involved in their production, conversion and clearance (PubMed:<a href="http://www.uniprot.org/citations/6860692" target="\_blank">6860692</a>, PubMed:<a href="http://www.uniprot.org/citations/2762297" target="\_blank">2762297</a>, PubMed:<a href="http://www.uniprot.org/citations/1911868" target="\_blank">1911868</a>, PubMed:<a href="http://www.uniprot.org/citations/1917954" target="\_blank">1917954</a>, PubMed:<a href="http://www.uniprot.org/citations/9395455" target="\_blank">9395455</a>, PubMed:<a href="http://www.uniprot.org/citations/14754908" target="\_blank">14754908</a>, PubMed:<a href="http://www.uniprot.org/citations/23620513" target="\_blank">23620513</a>). Apolipoproteins are amphipathic molecules that interact both with lipids of the lipoprotein particle core and the aqueous environment of the plasma (PubMed:<a href="http://www.uniprot.org/citations/6860692" target="\_blank">6860692</a>, PubMed:<a href="http://www.uniprot.org/citations/2762297" target="\_blank">2762297</a>, PubMed:<a href="http://www.uniprot.org/citations/9395455" target="\_blank">9395455</a>). As such, APOE associates with chylomicrons, chylomicron remnants, very low density lipoproteins (VLDL) and intermediate density lipoproteins (IDL) but shows a preferential binding to high-density lipoproteins (HDL) (PubMed:<a href="http://www.uniprot.org/citations/6860692" target="\_blank">6860692</a>, PubMed:<a href="http://www.uniprot.org/citations/1911868" target="\_blank">1911868</a>). It also binds a wide range of cellular receptors including the LDL receptor/LDLR, the LDL receptor-related proteins LRP1, LRP2 and LRP8 and the very low-density lipoprotein receptor/VLDLR that mediate the cellular uptake of the APOE-containing lipoprotein particles (PubMed:<a href="http://www.uniprot.org/citations/2762297" target="\_blank">2762297</a>, PubMed:<a href="http://www.uniprot.org/citations/1917954" target="\_blank">1917954</a>, PubMed:<a href="http://www.uniprot.org/citations/7768901" target="\_blank">7768901</a>, PubMed:<a href="http://www.uniprot.org/citations/8939961" target="\_blank">8939961</a>, PubMed:<a href="http://www.uniprot.org/citations/12950167" target="\_blank">12950167</a>, PubMed:<a href="http://www.uniprot.org/citations/20030366" target="\_blank">20030366</a>, PubMed:<a href="http://www.uniprot.org/citations/2063194" target="\_blank">2063194</a>, PubMed:<a href="http://www.uniprot.org/citations/8756331" target="\_blank">8756331</a>, PubMed:<a href="http://www.uniprot.org/citations/20303980" target="\_blank">20303980</a>, PubMed:<a href="http://www.uniprot.org/citations/1530612" target="\_blank">1530612</a>, PubMed:<a href="http://www.uniprot.org/citations/7635945" target="\_blank">7635945</a>). Finally, APOE has also a heparin-binding activity and binds heparan- sulfate proteoglycans on the surface of cells, a property that supports the capture and the receptor-mediated uptake of APOE-containing lipoproteins by cells (PubMed:<a href="http://www.uniprot.org/citations/9395455" target="\_blank">9395455</a>, PubMed:<a href="http://www.uniprot.org/citations/9488694" target="\_blank">9488694</a>, PubMed:<a href="http://www.uniprot.org/citations/23676495" target="\_blank">23676495</a>, PubMed:<a href="http://www.uniprot.org/citations/7635945" target="\_blank">7635945</a>). A main function of APOE is to mediate lipoprotein clearance through the uptake of chylomicrons, VLDLs, and HDLs

by hepatocytes (PubMed:<a href="http://www.uniprot.org/citations/1911868" target="\_blank">1911868</a>, PubMed:<a href="http://www.uniprot.org/citations/1917954" target="\_blank">1917954</a>, PubMed:<a href="http://www.uniprot.org/citations/9395455" target="\_blank">9395455</a>, PubMed:<a href="http://www.uniprot.org/citations/23676495" target="\_blank">23676495</a>, PubMed:<a href="http://www.uniprot.org/citations/29516132" target="\_blank">29516132</a>). APOE is also involved in the biosynthesis by the liver of VLDLs as well as their uptake by peripheral tissues ensuring the delivery of triglycerides and energy storage in muscle, heart and adipose tissues (PubMed:<a href="http://www.uniprot.org/citations/2762297" target="\_blank">2762297</a>, PubMed:<a href="http://www.uniprot.org/citations/29516132" target="\_blank">29516132</a>). By participating in the lipoprotein-mediated distribution of lipids among tissues, APOE plays a critical role in plasma and tissues lipid homeostasis (PubMed:<a href="http://www.uniprot.org/citations/2762297" target="\_blank">2762297</a>, PubMed:<a href="http://www.uniprot.org/citations/1917954" target="\_blank">1917954</a>, PubMed:<a href="http://www.uniprot.org/citations/29516132" target="\_blank">29516132</a>). APOE is also involved in two steps of reverse cholesterol transport, the HDLs-mediated transport of cholesterol from peripheral tissues to the liver, and thereby plays an important role in cholesterol homeostasis (PubMed:<a href="http://www.uniprot.org/citations/9395455" target="\_blank">9395455</a>, PubMed:<a href="http://www.uniprot.org/citations/14754908" target="\_blank">14754908</a>, PubMed:<a href="http://www.uniprot.org/citations/23620513" target="\_blank">23620513</a>). First, it is functionally associated with ABCA1 in the biogenesis of HDLs in tissues (PubMed:<a href="http://www.uniprot.org/citations/14754908" target="\_blank">14754908</a>, PubMed:<a href="http://www.uniprot.org/citations/23620513" target="\_blank">23620513</a>). Second, it is enriched in circulating HDLs and mediates their uptake by hepatocytes (PubMed:<a href="http://www.uniprot.org/citations/9395455" target="\_blank">9395455</a>). APOE also plays an important role in lipid transport in the central nervous system, regulating neuron survival and sprouting (PubMed:<a href="http://www.uniprot.org/citations/8939961" target="\_blank">8939961</a>, PubMed:<a href="http://www.uniprot.org/citations/25173806" target="\_blank">25173806</a>). APOE is also involved in innate and adaptive immune responses, controlling for instance the survival of myeloid-derived suppressor cells (By similarity). Binds to the immune cell receptor LILRB4 (PubMed:<a href="http://www.uniprot.org/citations/30333625" target="\_blank">30333625</a>). APOE may also play a role in transcription regulation through a receptor-dependent and cholesterol-independent mechanism, that activates MAP3K12 and a non-canonical MAPK signal transduction pathway that results in enhanced AP-1-mediated transcription of APP (PubMed:<a href="http://www.uniprot.org/citations/28111074" target="\_blank">28111074</a>).

### Cellular Location

Secreted. Secreted, extracellular space. Secreted, extracellular space, extracellular matrix. Extracellular vesicle. Endosome, multivesicular body. Note=In the plasma, APOE is associated with chylomicrons, chylomicrons remnants, VLDL, LDL and HDL lipoproteins (PubMed:1911868, PubMed:8340399). Lipid poor oligomeric APOE is associated with the extracellular matrix in a calcium- and heparan-sulfate proteoglycans-dependent manner (PubMed:9488694) Lipidation induces the release from the extracellular matrix (PubMed:9488694). Colocalizes with CD63 and PMEL at exosomes and in intraluminal vesicles within multivesicular endosomes

### Tissue Location

Produced by several tissues and cell types and mainly found associated with lipid particles in the plasma, the interstitial fluid and lymph (PubMed:25173806). Mainly synthesized by liver hepatocytes (PubMed:25173806). Significant quantities are also produced in brain, mainly by astrocytes and glial cells in the cerebral cortex, but also by neurons in frontal cortex and hippocampus (PubMed:3115992, PubMed:10027417). It is also expressed by cells of the peripheral nervous system (PubMed:10027417, PubMed:25173806). Also expressed by adrenal gland, testis, ovary, skin, kidney, spleen and adipose tissue and macrophages in various tissues (PubMed:25173806)

## **ApoE3 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **ApoE3 Antibody - Images**

## **ApoE3 Antibody - Background**

ApoE belongs to a group of proteins that bind reversibly with lipoprotein and play an important role in lipid metabolism. ApoE is an important constituent of all plasma lipoproteins. It's interaction with specific ApoE receptor enables uptake of chylomicron remnants by liver cells, which is an essential step during normal lipid metabolism. It also binds with the LDL receptor (apo B/E). Defects in ApoE are a cause of hyperlipoproteinemia type III. ApoE exists in three major isoforms; E2, E3, and E4, which differ from one another by a single amino-acid substitution. E3 is the most common isoform and is present in 40-90% of the population.