

Wnt-1 Antibody
Rabbit Polyclonal Antibody
Catalog # ABV11060**Specification**

Wnt-1 Antibody - Product Information

| | |
|-------------------|------------------------|
| Application | WB |
| Primary Accession | P04628 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Calculated MW | 40982 |

Wnt-1 Antibody - Additional Information**Gene ID 7471**

| | |
|---------------------|---|
| Application & Usage | Western blot analysis (0.5-4 µg/ml). However, the optimal conditions should be determined individually. Recombinant human Wnt-1 can be used as a positive control. |
|---------------------|---|

Other Names

wnt1, wnt 1, wnt-1, INT 1 , INT1, wingless type MMTV integration site family member 1

Target/Specificity

Wnt-1

Antibody Form

Liquid

Appearance

Colorless liquid

Formulation

100 µg (0.5 mg/ml) affinity purified rabbit anti-human Wnt-1 polyclonal antibody in phosphate buffered saline (PBS), pH 7.2, containing 30% glycerol, 0.5% BSA, and 0.01% thimerosal.

Handling

The antibody solution should be gently mixed before use.

Reconstitution & Storage

-20 °C

Background Descriptions**Precautions**

Wnt-1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Wnt-1 Antibody - Protein Information

Name WNT1

Synonyms INT1

Function

Ligand for members of the frizzled family of seven transmembrane receptors (Probable). Acts in the canonical Wnt signaling pathway by promoting beta-catenin-dependent transcriptional activation (PubMed:23499309, PubMed:26902720, PubMed:28528193, PubMed:23656646). In some developmental processes, is also a ligand for the coreceptor RYK, thus triggering Wnt signaling (By similarity). Plays an essential role in the development of the embryonic brain and central nervous system (CNS) (By similarity). Has a role in osteoblast function, bone development and bone homeostasis (PubMed:23499309, PubMed:23656646).

Cellular Location

Secreted, extracellular space, extracellular matrix. Secreted

Wnt-1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Wnt-1 Antibody - Images

Wnt-1 Antibody - Background

Wnt-1 is a secreted protein that signals through the Frizzled family of cell surface receptors and is required for normal embryonic development. Wnt-1 activation induces a complex signaling cascade that ultimately leads to the increased expression of over fifty genes. An important component of Wnt-1 signaling is the stabilization, and resulting accumulation, of the intracellular signaling protein, beta-catenine. Wnt signaling induces and maintains the transformed phenotype and, in certain embryonic cell lines, supports self renewal in the absence of significant differentiation. Elevated levels of Wnt proteins are associated with tumorigenesis and are present in numerous human breast cancers. Mature human Wnt-1 is a glycosylated protein containing 343 amino acid residues.