

K2/spice, Synthetic Cannabinoids Antibody

Rabbit Polyclonal Antibody Catalog # ABV11169

Specification

K2/spice, Synthetic Cannabinoids Antibody - Product Information

Application Reactivity Host Clonality Isotype

E All Species Rabbit Polyclonal Rabbit IgG

K2/spice, Synthetic Cannabinoids Antibody - Additional Information

Application & Usage

Other Names CB, CX5 ELISA (for 96-well plate coating use 1-3 μ g/mL).

Target/Specificity K2/spice

Antibody Form Liquid

Appearance Colorless liquid

Formulation 2 mg/ml of rabbit IgG in phosphate buffered saline with 0.05% sodium azide preservative.

Handling The antibody solution should be gently mixed before use.

Reconstitution & Storage -20 °C

Background Descriptions

Precautions K2/spice, Synthetic Cannabinoids Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

K2/spice, Synthetic Cannabinoids Antibody - Protein Information



K2/spice, Synthetic Cannabinoids Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

K2/spice, Synthetic Cannabinoids Antibody - Images

K2/spice, Synthetic Cannabinoids Antibody - Background

Cannabinoids are a class of diverse chemical compounds that activate cannabinoid receptors on cells that repress neurotransmitter release in the brain. They are active chemicals in Cannabis that cause drug-like effects throughout the body, including the central nervous system and the immune system. Anti-K2/spice (synthetic cannabinoids), is a rabbit polyclonal IgG antibody. It has been used in a competitive ELISA format to test the presence of JWH-018, JWH-073, JWH-122, JWH-019, JWH-081, AM-2201 and related compounds and their metabolites in samples such as urine, whole blood, serum, and plasma (see Arntson et al, 2013). Note: If this antibody is used in an immunoassay to detect synthetic cannabinoids, suspect test samples must be confirmed using an alternative analytical method, for example LC-MS-MS.