Western blot: HeLa cells, ELISA: Peptides.

Western Blot: 1:500, ELISA: 1:100 - 1:500.



JMJD2a polyclonal antibody

Rabbit Polyclonal Antibody Catalog # ABV11385

Specification

JMJD2a polyclonal antibody - Product Information

Application WB, E
Primary Accession O75164
Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 120662

JMJD2a polyclonal antibody - Additional Information

Gene ID 9682

Positive Control Application & Usage **Other Names** JHDM3A, KDM4A, JMJD2

Target/Specificity JMJD2a

Antibody Form Liquid

Appearance Colorless liquid

Formulation

In PBS with 0.05% sodium azide and 0.05% ProClin 300.

Handling

The antibody solution should be gently mixed before use.

Reconstitution & Storage -20 °C

Background Descriptions

Precautions

JMJD2a polyclonal antibody is for research use only and not for use in diagnostic or therapeutic procedures.

JMJD2a polyclonal antibody - Protein Information



Name KDM4A

Synonyms JHDM3A, JMJD2, JMJD2A, KIAA0677

Function

Histone demethylase that specifically demethylates 'Lys-9' and 'Lys-36' residues of histone H3, thereby playing a central role in histone code (PubMed:26741168). Does not demethylate histone H3 'Lys- 4', H3 'Lys-27' nor H4 'Lys-20'. Demethylates trimethylated H3 'Lys-9' and H3 'Lys-36' residue, while it has no activity on mono- and dimethylated residues. Demethylation of Lys residue generates formaldehyde and succinate. Participates in transcriptional repression of ASCL2 and E2F-responsive promoters via the recruitment of histone deacetylases and NCOR1, respectively.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00537, ECO:0000269|PubMed:15927959, ECO:0000269|PubMed:16024779}

Tissue Location Ubiquitous..

JMJD2a polyclonal antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

JMJD2a polyclonal antibody - Images

JMJD2a polyclonal antibody - Background

JMJD2a belongs to the JMJD2 family of histone demethylases which play an important role in the establishment of the histone code. JMJD2a specifically demethylates the trimethylated K9 and K36 of histone H3, thereby converting these lysines to the dimethylated form. It has no activity towards H3K4, H3K27 and H4K20, or to the mono- and dimethylated H3K9 and H3K36. JMJD2a plays a role in the transcriptional repression of ASCL2 and E2F-responsive promoters via the recruitment of histone deacetylases and NCOR1, respectively.