

### **PTP1B Antibody**

Purified Rabbit Polyclonal Antibody Catalog # ABV11509

# Specification

# PTP1B Antibody - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Isotype
Calculated MW

WB <u>P18031</u> Human Rabbit Polyclonal Rabbit IgG 49967

### **PTP1B** Antibody - Additional Information

Gene ID 5770

**Other Names** Tyrosine-protein phosphatase non-receptor type 1, Protein-tyrosine phosphatase 1B, PTP-1B, PTP1B

Target/Specificity PTP1B

Formulation 100  $\mu$ g (0.25 mg/ml) purified rabbit Ig polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Handling The antibody solution should be gently mixed before use.

**Background Descriptions** 

**Precautions** PTP1B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### **PTP1B Antibody - Protein Information**

Name PTPN1

Synonyms PTP1B

#### Function

Tyrosine-protein phosphatase which acts as a regulator of endoplasmic reticulum unfolded protein response. Mediates dephosphorylation of EIF2AK3/PERK; inactivating the protein kinase activity of EIF2AK3/PERK. May play an important role in CKII- and p60c- src-induced signal transduction



cascades. May regulate the EFNA5-EPHA3 signaling pathway which modulates cell reorganization and cell-cell repulsion. May also regulate the hepatocyte growth factor receptor signaling pathway through dephosphorylation of MET.

**Cellular Location** 

Endoplasmic reticulum membrane; Peripheral membrane protein; Cytoplasmic side Note=Interacts with EPHA3 at the cell membrane


#### **Tissue Location**

Expressed in keratinocytes (at protein level).

# PTP1B Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>
- **PTP1B Antibody Images**



## PTP1B Antibody - Background

PTP1B is the founding member of the protein tyrosine phosphatase (PTP) family, which was isolated and identified based on its enzymatic activity and amino acid sequence. PTPs catalyze the hydrolysis of the phosphate monoesters specifically on tyrosine residues. Members of the PTP family share a highly conserved catalytic motif, which is essential for the catalytic activity. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP has been shown to act as a negative regulator of insulin signaling by dephosphorylating the phosphotryosine residues of insulin receptor kinase. This PTP was also reported to dephosphorylate epidermal growth factor receptor kinase, as well as JAK2 and TYK2 kinases, which implicated the role of this PTP in cell growth



control, and cell response to interferon stimulation.