

## **TAOK1** Antibody

Purified Rabbit Polyclonal Antibody Catalog # ABV11558

## **Specification**

## **TAOK1 Antibody - Product Information**

Application
Primary Accession
Other Accession
Reactivity
Host
Clonality
Isotype
Calculated MW

WB
088664
EDM05292
Human, Mouse, Rat
Rabbit
Polyclonal
Rabbit IgG
115952

### **TAOK1** Antibody - Additional Information

#### Gene ID 286993

### **Other Names**

TAOK1, TAO kinase 1, MARKK, Microtubule affinity regulating kinase kinase, MAP3K16, PSK2, TAO1

# **Target/Specificity**

TAOK1

#### **Formulation**

 $100 \mu g$  (0.5mg/ml) affinity purified rabbit polyclonal antibody in phosphate-buffered saline (PBS) containing 30% glycerol, 0.5% BSA and 0.01% thimerosal.

#### Handling

The antibody solution should be gently mixed before use.

#### **Background Descriptions**

#### **Precautions**

TAOK1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## **TAOK1 Antibody - Protein Information**

#### Name Taok1

## Synonyms Tao1

## **Function**

Serine/threonine-protein kinase involved in various processes such as p38/MAPK14 stress-activated MAPK cascade, DNA damage response and regulation of cytoskeleton stability. Phosphorylates MAP2K3, MAP2K6 and MARK2. Acts as an activator of the p38/MAPK14



stress-activated MAPK cascade by mediating phosphorylation and subsequent activation of the upstream MAP2K3 and MAP2K6 kinases. Involved in G-protein coupled receptor signaling to p38/MAPK14. In response to DNA damage, involved in the G2/M transition DNA damage checkpoint by activating the p38/MAPK14 stress-activated MAPK cascade, probably by mediating phosphorylation of MAP2K3 and MAP2K6. Acts as a regulator of cytoskeleton stability by phosphorylating 'Thr-208' of MARK2, leading to activate MARK2 kinase activity and subsequent phosphorylation and detachment of MAPT/TAU from microtubules. Also acts as a regulator of apoptosis: regulates apoptotic morphological changes, including cell contraction, membrane blebbing and apoptotic bodies formation via activation of the MAPK8/JNK cascade. During fetal development, it plays an essential role in the regulation of neuronal differentiation and migration to the cortical plate (By similarity).


**Cellular Location** Cytoplasm.

## **TAOK1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## **TAOK1 Antibody - Images**



#### **TAOK1 Antibody - Background**

Serine/threonine-protein kinase TAO1 is an enzyme that in humans is encoded by the TAOK1 gene