

**Anti-Lactoferrin Antibody (10D04)**  
**Mouse Monoclonal Antibody**  
**Catalog # ABV12107****Specification**

---

**Anti-Lactoferrin Antibody (10D04) - Product Information**

Application	E
Primary Accession	<a href="#">P02788</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse IgG1, κ

**Anti-Lactoferrin Antibody (10D04) - Additional Information****Gene ID** 4057**Positive Control** ELISA**Other Names**

Growth-inhibiting protein 12, Talalactoferrin, Lactotransferrin

**Target/Specificity**

Lactoferrin

**Antibody Form**

Liquid

**Appearance**

Colorless liquid

**Reconstitution & Storage**

-20 °C

**Background Descriptions****Precautions**

Anti-Lactoferrin Antibody (10D04) is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-Lactoferrin Antibody (10D04) - Protein Information****Name** LTF ([HGNC:6720](#))**Synonyms** GIG12, LF**Function**

Transferrins are iron binding transport proteins which can bind two Fe(3+) ions in association with

the binding of an anion, usually bicarbonate.

### Cellular Location

[Isoform 1]: Secreted. Cytoplasmic granule. Note=Secreted into most exocrine fluids by various endothelial cells Stored in the secondary granules of neutrophils

### Tissue Location


High levels are found in saliva and tears, intermediate levels in serum and plasma, and low levels in urine. In kidney, detected in the distal collecting tubules in the medulla but not in the cortical region or in blood vessels. Detected in peripheral blood neutrophils (at protein level). Isoform 1 and isoform DeltaLf are expressed in breast, prostate, spleen, pancreas, kidney, small intestine, lung, skeletal muscle, uterus, thymus and fetal liver Isoform 1 is expressed in brain, testis and peripheral blood leukocytes; isoform DeltaLf is barely detectable in these tissues Isoform DeltaLf is expressed in placenta, liver and ovary; isoform 1 is barely detectable in these tissues. In kidney, isoform 1 is expressed at high levels in the collecting tubules of the medulla but at very low levels in the cortex.

## Anti-Lactoferrin Antibody (10D04) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-Lactoferrin Antibody (10D04) - Images



The antibodies were evaluated for their ability to work as matched pairs in a mono-mono sandwich ELISA format. Biotin conjugates of both antibodies were prepared for use as tracers. The antibodies demonstrated the ability to work in both capture/tracer configurations

## Anti-Lactoferrin Antibody (10D04) - Background

Lactoferrin is a glycoprotein present in exocrine secretions and in the secondary granules of polymorphonuclear neutrophils. It is the iron binding protein in milk. Lactoferrin is known to regulate intestinal iron absorption, and to participate in the defense against bacteria. Lactoferrin is part of the transferrin family of proteins, an iron binding glycoprotein involved in host defense

against infection and is a modulator of inflammatory reactions.