

**CD3**

**Rabbit Monoclonal antibody(Mab)**  
**Catalog # AD80004**

**Specification****CD3 - Product info**

Application	IHC-P
Primary Accession	<a href="#">P04234</a>
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	18930

**CD3 - Additional info**

Gene ID	915
Gene Name	CD3D
<b>Other Names</b>	
T-cell surface glycoprotein CD3 delta chain, T-cell receptor T3 delta chain, CD3d, CD3D, T3D	
<b>Dilution</b>	
IHC-P~Ready-to-use	
<b>Storage</b>	
Maintain refrigerated at 2-8°C	
Precautions	<b>CD3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.</b>

**CD3 - Protein Information****Name** CD3D

Synonyms	T3D
Function	Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in

the activation of downstream signaling pathways (PubMed:[2470098](#)). In addition of this role of signal transduction in T-cell activation, CD3D plays an essential role in thymocyte differentiation. Indeed, participates in correct intracellular TCR-CD3 complex assembly and surface expression. In absence of a functional TCR-CD3 complex, thymocytes are unable to differentiate properly. Interacts with CD4 and CD8 and thus serves to establish a functional link between the TCR and coreceptors CD4 and CD8, which is needed for activation and positive selection of CD4 or CD8 T-cells(PubMed:[12215456](#)).

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location


CD3D is mostly present on T-lymphocytes with its TCR-CD3 partners. Present also in fetal NK-cells

## CD3 - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## CD3 - Images



Tonsil



Immunohistochemical analysis of paraffin-embedded human tonsil tissue using AD80004 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a Citrate buffer (pH 6.0). Samples were incubated with primary antibody(Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems Abcepta:AR005 was used as the secondary antibody.