

СК20

Rabbit Monoclonal antibody(Mab) Catalog # AD80024

Specification

CK20 - Product info

Application Primary Accession Reactivity Host Clonality Calculated MW

P35900 Human Rabbit Monoclonal 48487

IHC-P

CK20 - Additional info

Gene ID 54474 Gene Name KRT20 Other Names Keratin, type I cytoskeletal 20, Cytokeratin-20, CK-20, Keratin-20, K20, Protein IT, KRT20

Dilution IHC-P~~Ready-to-use

Storage Maintain refrigerated at 2-8°C

Precautions

CK20 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CK20 - Protein Information

Name KRT20

Function

Cellular Location Tissue Location Plays a significant role in maintaining keratin filament organization in intestinal epithelia. When phosphorylated, plays a role in the secretion of mucin in the small intestine (By similarity). Cytoplasm Expressed predominantly in the intestinal epithelium. Expressed in luminal cells of colonic mucosa. Also expressed in the Merkel cells of keratinized oral mucosa; specifically at the tips of some rete ridges of the gingival mucosa, in the basal layer of the palatal mucosa.



CK20 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>
- CK20 Images



Colon cancer



Immunohistochemical analysis of paraffin-embedded colorectal carcinoma; tissue using AD80024 performed on the Abcarta® FAIP-30 Fully automated IHC platform.Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a Citrate buffer (pH6. 0).Samples were incubated with primary antibody(Ready-to-use) for 15 min at room temperature. AmpSeeTM Detection Systems[Abcepta:AR005] was used as the secondary antibody.