

Vimentin
Mouse Monoclonal antibody(Mab)
Catalog # AD80040**Specification****Vimentin - Product info**

Application	IHC-P
Primary Accession	P08670
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	53652

Vimentin - Additional info

Gene ID	7431
Gene Name	VIM
Other Names	
Vimentin, VIM	

Dilution
IHC-P~ Ready-to-use**Storage**
Maintain refrigerated at 2-8°C**Precautions**
Vimentin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.**Vimentin - Protein Information****Name** VIM ([HGNC:12692](#))


Function	Vimentins are class-III intermediate filaments found in various non-epithelial cells, especially mesenchymal cells. Vimentin is attached to the nucleus, endoplasmic reticulum, and mitochondria, either laterally or terminally.
Cellular Location	Cytoplasm. Cytoplasm, cytoskeleton. Nucleus matrix
Tissue Location	{ECO:0000250 UniProtKB:P31000} Highly expressed in fibroblasts, some expression in T- and B-lymphocytes, and little or no expression in Burkitt's lymphoma cell lines. Expressed in many hormone- independent mammary carcinoma cell lines

Vimentin - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Vimentin - Images



Liver



Immunohistochemical analysis of paraffin-embedded gastrointestinal stromal tumor tissue using AD80305 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a Citrate buffer (pH 6.0). Samples were incubated with primary antibody(Ready-to-use) for 15 min at room temperature. AmpSeeTM Detection Systems [Abcepta:AR005] was used as the secondary antibody.