

## MUC2

Mouse Monoclonal antibody(Mab) Catalog # AD80096

### **Specification**

## MUC2 - Product info

Application Primary Accession Reactivity Host Clonality Calculated MW IHC-P <u>002817</u> Human Mouse Monoclonal 550850

# MUC2 - Additional info

Gene Name MUC2 Other Names Mucin-2, MUC-2, Intestinal mucin-2, MUC2 {ECO:0000303|PubMed:8300571, ECO:0000312|HGNC:HGNC:7512}

Dilution IHC-P~~Ready-to-use

Storage Maintain refrigerated at 2-8°C

Precautions

MUC2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### **MUC2 - Protein Information**

Name MUC2 {ECO:0000303|PubMed:8300571, ECO:0000312|HGNC:HGNC:7512}

| Synonyms<br>Function | SMUC<br>Coats the epithelia of the intestines,<br>airways, and other mucus<br>membrane-containing organs. Thought to<br>provide a protective, lubricating barrier<br>against particles and infectious agents at<br>mucosal surfaces. Major constituent of<br>both the inner and outer mucus layers of<br>the colon and may play a role in excluding<br>bacteria from the inner mucus layer. |
|----------------------|---|
| Cellular Location    | Secreted. Note=In the intestine, secreted into the inner and outer mucus layers.  |
| Tissue Location      | Colon, small intestine, colonic tumors,<br>bronchus, cervix and gall bladder  |



## MUC2 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>
- **MUC2** Images



Colon cancer



Immunohistochemical analysis of paraffin-embedded colorectal carcinoma; tissue using AD80096 performed on the Abcarta® FAIP-30 Fully automated IHC platform.Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody(Ready-to-use) for 15 min at room temperature. AmpSeeTM Detection Systems[Abcepta:AR005] was used as the secondary antibody.