

**CFTR**  
**Mouse Monoclonal antibody(Mab)**  
**Catalog # AD80156****Specification**

---

**CFTR - Product info**

Application	IHC-P, IHC
Primary Accession	<a href="#">P13569</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	168142

**CFTR - Additional info**

Gene ID	1080
Gene Name	CFTR

**Other Names**

Cystic fibrosis transmembrane conductance regulator, CFTR, ATP-binding cassette sub-family C member 7, Channel conductance-controlling ATPase, 5.6.1.6, cAMP-dependent chloride channel, CFTR, ABCC7

**Dilution**

IHC-P~~Ready-to-use  
IHC~~Ready-to-use

Storage	This product is stored at 2-95 °C, please use it within the expiration date.
Precautions	CFTR Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**CFTR - Protein Information****Name** CFTR

Synonyms	ABCC7
Function	Epithelial ion channel that plays an important role in the regulation of epithelial ion and water transport and fluid homeostasis (PubMed: <a href="#">26823428</a> ). Mediates the transport of chloride ions across the cell membrane (PubMed: <a href="#">10792060</a> , PubMed: <a href="#">11524016</a> , PubMed: <a href="#">11707463</a> , PubMed: <a href="#">12519745</a> , PubMed: <a href="#">15010471</a> , PubMed: <a href="#">12588899</a> , PubMed: <a href="#">17036051</a> , PubMed: <a href="#">19398555</a> , PubMed: <a href="#">19621064</a> , PubMed: <a href="#">22178883</a> , PubMed: <a href="#">25330774</a> , PubMed: <a href="#">1712898</a> , PubMed: <a href="#">8910473</a> ,

## Cellular Location

PubMed:[9804160](#), PubMed:[12529365](#), PubMed:[17182731](#), PubMed:[26846474](#), PubMed:[28087700](#)). Channel activity is coupled to ATP hydrolysis (PubMed:[8910473](#)). The ion channel is also permeable to  $\text{HCO}_3^-$ ; selectivity depends on the extracellular chloride concentration (PubMed:[15010471](#), PubMed:[19019741](#)). Exerts its function also by modulating the activity of other ion channels and transporters (PubMed:[12403779](#), PubMed:[22178883](#), PubMed:[22121115](#), PubMed:[27941075](#)). Plays an important role in airway fluid homeostasis (PubMed:[16645176](#), PubMed:[19621064](#), PubMed:[26823428](#)). Contributes to the regulation of the pH and the ion content of the airway surface fluid layer and thereby plays an important role in defense against pathogens (PubMed:[14668433](#), PubMed:[16645176](#), PubMed:[26823428](#)). Modulates the activity of the epithelial sodium channel (ENaC) complex, in part by regulating the cell surface expression of the ENaC complex (PubMed:[17434346](#), PubMed:[27941075](#), PubMed:[17182731](#)). Inhibits the activity of the ENaC channel containing subunits SCNN1A, SCNN1B and SCNN1G (PubMed:[17182731](#)). Inhibits the activity of the ENaC channel containing subunits SCNN1D, SCNN1B and SCNN1G, but not of the ENaC channel containing subunits SCNN1A, SCNN1B and SCNN1G (PubMed:[27941075](#)). May regulate bicarbonate secretion and salvage in epithelial cells by regulating the transporter SLC4A7 (PubMed:[12403779](#)). Can inhibit the chloride channel activity of ANO1 (PubMed:[22178883](#)). Plays a role in the chloride and bicarbonate homeostasis during sperm epididymal maturation and capacitation (PubMed:[19923167](#), PubMed:[27714810](#)).

Apical cell membrane; Multi-pass membrane protein {ECO:0000269|Ref.54}. Early endosome membrane; Multi-pass membrane protein {ECO:0000269|Ref.54}. Cell membrane; Multi-pass membrane protein {ECO:0000269|Ref.54}. Recycling endosome membrane; Multi-pass membrane protein {ECO:0000269|Ref.54}. Endoplasmic reticulum membrane; Multi-pass membrane protein {ECO:0000269|Ref.54}. Nucleus {ECO:0000250|UniProtKB:P34158}. Note=The channel is internalized from the cell surface into an endosomal recycling

#### Tissue Location


compartment, from where it is recycled to the cell membrane (PubMed:17462998, PubMed:19398555, PubMed:20008117). In the oviduct and bronchus, detected on the apical side of epithelial cells, but not associated with cilia (PubMed:22207244). In Sertoli cells, a processed product is detected in the nucleus (By similarity). ER stress induces GORASP2-mediated unconventional (ER/Golgi- independent) trafficking of core-glycosylated CFTR to cell membrane (PubMed:21884936). {ECO:0000250|UniProtKB:P34158, ECO:0000269|PubMed:19398555, ECO:0000269|PubMed:20008117, ECO:0000269|PubMed:21884936, ECO:0000269|PubMed:22207244, ECO:0000305|PubMed:17462998} Expressed in the respiratory airway, including bronchial epithelium, and in the female reproductive tract, including oviduct (at protein level) (PubMed:22207244, PubMed:15716351). Detected in pancreatic intercalated ducts in the exocrine tissue, on epithelial cells in intralobular striated ducts in sublingual salivary glands, on apical membranes of crypt cells throughout the small and large intestine, and on the reabsorptive duct in eccrine sweat glands (PubMed:1284548, PubMed:28130590). Detected on the equatorial segment of the sperm head (at protein level) (PubMed:19923167). Detected in nasal and bronchial superficial epithelium (PubMed:15716351). Expressed by the central cells on the sebaceous glands, dermal adipocytes and, at lower levels, by epithelial cells (PubMed:28130590)


#### CFTR - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### CFTR - Images





Pancreas



Immunohistochemical analysis of paraffin-embedded human pancreas tissue using AD80156 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9.0). Samples were incubated with primary antibody (Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems [Abcepta:AR005] was used as the secondary antibody.