

**TPO**  
**Mouse Monoclonal antibody(Mab)**  
**Catalog # AD80232**

## Specification

---

### TPO - Product info

Application	<b>IHC-P</b>
Primary Accession	<a href="#">P07202</a>
Reactivity	<b>Human</b>
Host	<b>Mouse</b>
Clonality	<b>Monoclonal</b>
Calculated MW	<b>102963</b>

### TPO - Additional info

Gene ID	<b>7173</b>
Gene Name	<b>TPO</b> ( <a href="#">HGNC:12015</a> )

#### Other Names

Thyroid peroxidase, TPO, 1.11.1.8, TPO (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=12015" target="\_blank">HGNC:12015</a>)

#### Dilution

IHC-P~~Ready-to-use

#### Storage

Maintain refrigerated at 2-8°C

#### Precautions

**TPO Antibody is for research use only and not for use in diagnostic or therapeutic procedures.**

### TPO - Protein Information

**Name** TPO ([HGNC:12015](#))

#### Function

**Iodination and coupling of the hormonogenic tyrosines in thyroglobulin to yield the thyroid hormones T(3) and T(4). Membrane; Single-pass type I membrane protein**

#### Cellular Location

### TPO - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)




- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **TPO - Images**



Papillary thyroid carcinoma,



Immunohistochemical analysis of paraffin-embedded prostatic cancer tissue using AD80295 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9.0). Samples were incubated with primary antibody (Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems [Abcepta:AR005] was used as the secondary antibody.