



CD38

Rabbit Monoclonal antibody(Mab)
Catalog # AD80356

Specification

CD38 - Product info

Application IHC-P
Primary Accession P28907
Reactivity Human
Host Rabbit
Clonality Monoclonal
Calculated MW 34328

CD38 - Additional info

Gene ID 952 Gene Name CD38

Other Names

ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1, 3.2.2.-, 3.2.2.6, 2'-phospho-ADP-ribosyl cyclase, 2'-phospho-ADP-ribosyl cyclase/2'-phospho-cyclic-ADP-ribose transferase, 2.4.99.20, 2'-phospho-cyclic-ADP-ribose transferase, ADP-ribosyl cyclase 1, ADPRC 1, Cyclic ADP-ribose hydrolase 1, cADPR hydrolase 1, T10, CD38, CD38

Dilution

IHC-P~~Ready-to-use

Storage

Maintain refrigerated at 2-8°C

Precautions CD38 Antibody is for research use only and

not for use in diagnostic or therapeutic

procedures.

CD38 - Protein Information

Name CD38

Function Synthesizes the second messagers cyclic

ADP-ribose and nicotinate-adenine dinucleotide phosphate, the former a second messenger for glucose-induced insulin secretion. Also has cADPr hydrolase activity. Also moonlights as a receptor in

cells of the immune system.

Cellular Location Membrane; Single-pass type II membrane

protein

Tissue Location Expressed at high levels in pancreas, liver,

Page 1/2

kidney, brain, testis, ovary, placenta, malignant lymphoma and neuroblastoma.




CD38 - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

CD38 - Images



Tonsil



Immunohistochemical analysis of paraffin-embedded human tonsil tissue using AD80356 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody (Ready-to-use) for 15 min at room temperature. AmpSeeTM Detection Systems Abcepta: AR005 was used as the secondary antibody.