

**LAG3**  
**Rabbit Monoclonal antibody(Mab)**  
**Catalog # AD80600**

## Specification

---

### LAG3 - Product info

Application	IHC-P
Primary Accession	<a href="#">P18627</a>
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	57449

### LAG3 - Additional info

Gene ID **3902**

#### Other Names

Lymphocyte activation gene 3 protein, LAG-3, CD223, Secreted lymphocyte activation gene 3 protein, sLAG-3, LAG3 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=6476](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=6476)), FDC

#### Dilution

IHC-P~~N/A

#### Storage

Maintain refrigerated at 2-8°C

### LAG3 - Protein Information

Name LAG3 ([HGNC:6476](#))

#### Synonyms

#### Function

#### FDC

Lymphocyte activation gene 3 protein: Inhibitory receptor on antigen activated T-cells (PubMed:[20421648](#), PubMed:[7805750](#), PubMed:[8647185](#)). Delivers inhibitory signals upon binding to ligands, such as FGL1 (By similarity). FGL1 constitutes a major ligand of LAG3 and is responsible for LAG3 T-cell inhibitory function (By similarity). Following TCR engagement, LAG3 associates with CD3-TCR in the immunological synapse and directly inhibits T-cell activation (By similarity). May inhibit antigen-specific T-cell activation in synergy with PDCD1/PD-1, possibly by acting as a

	<b>coreceptor for PDCD1/PD-1 (By similarity). Negatively regulates the proliferation, activation, effector function and homeostasis of both CD8(+) and CD4(+) T-cells (PubMed:<a href="#">20421648</a>, PubMed:<a href="#">7805750</a>, PubMed:<a href="#">8647185</a>). Also mediates immune tolerance: constitutively expressed on a subset of regulatory T-cells (Tregs) and contributes to their suppressive function (By similarity). Also acts as a negative regulator of plasmacytoid dendritic cell (pDCs) activation (By similarity). Binds MHC class II (MHC-II); the precise role of MHC-II-binding is however unclear (PubMed:<a href="#">8647185</a>). [Lymphocyte activation gene 3 protein]: Cell membrane; Single-pass type I membrane protein</b>
Cellular Location	
Tissue Location	<b>Primarily expressed in activated T-cells and a subset of natural killer (NK) cells.</b>

### **LAG3 - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **LAG3 - Images**