

Phospho-Ezrin-T567 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AE1012b

Specification

Phospho-Ezrin-T567 Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Concentration Isotype Calculated MW WB <u>P15311</u> Human, Mouse, Rat Rabbit Polyclonal 1mg/ml Rabbit IgG 69413

Phospho-Ezrin-T567 Antibody - Additional Information

Gene ID 7430

Other Names Ezrin, Cytovillin, Villin-2, p81, EZR, VIL2

Target/Specificity

The antibody was affinity-purified from rabbit antiserum using epitope-specific phosphopeptide column, and the antibody against non-phosphopeptide was removed using non-phosphopeptide column corresponding to the phosphorylation site.

Dilution WB~~1:500~1:1000

Format affinity Purified IgG, in PBS, 0.02% sodium azide and 50% glycerol.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions Phospho-Ezrin-T567 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-Ezrin-T567 Antibody - Protein Information

Name EZR

Synonyms VIL2



Function

Probably involved in connections of major cytoskeletal structures to the plasma membrane. In epithelial cells, required for the formation of microvilli and membrane ruffles on the apical pole. Along with PLEKHG6, required for normal macropinocytosis.

Cellular Location

Apical cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection. Cell projection, microvillus membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, ruffle membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cell projection, microvillus {ECO:0000250|UniProtKB:P26040}. Note=Localization to the apical membrane of parietal cells depends on the interaction with PALS1 Localizes to cell extensions and peripheral processes of astrocytes (By similarity). Microvillar peripheral membrane protein (cytoplasmic side). {ECO:000250|UniProtKB:P31977}

Tissue Location

Expressed in cerebral cortex, basal ganglia, hippocampus, hypophysis, and optic nerve. Weakly expressed in brain stem and diencephalon. Stronger expression was detected in gray matter of frontal lobe compared to white matter (at protein level). Component of the microvilli of intestinal epithelial cells. Preferentially expressed in astrocytes of hippocampus, frontal cortex, thalamus, parahippocampal cortex, amygdala, insula, and corpus callosum. Not detected in neurons in most tissues studied

Phospho-Ezrin-T567 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Phospho-Ezrin-T567 Antibody - Images



Western blot analysis of extracts from A431 cells, untreated or EGFtreated (200ng/ml, 30min)



using Ezrin Antibody (T567) (#AE1012a, Lane 1 and 2) and Phospho-Ezrin-T567 Antibody (#AE1012b, Lane 3 and 4).

Phospho-Ezrin-T567 Antibody - Background

The cytoplasmic peripheral membrane protein encoded by this gene functions as a protein-tyrosine kinase substrate in microvilli. As a member of the ERM protein family, this protein serves as an intermediate between the plasma membrane and the actin cytoskeleton. This protein plays a key role in cell surface structure adhesion, migration and organization, and it has been implicated in various human cancers. A pseudogene located on chromosome 3 has been identified for this gene. Alternatively spliced variants have also been described for this gene.

Phospho-Ezrin-T567 Antibody - References

Ezrin immunoreactivity in renal cell carcinomas. Tuna B, et al. Anal Quant Cytol Histol, 2009 Oct. PMID 20701102.

Evaluation of candidate stromal epithelial cross-talk genes identifies association between risk of serous ovarian cancer and TERT, a cancer susceptibility hot-spot. Johnatty SE, et al. PLoS Genet, 2010 Jul 8. PMID 20628624.

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.

Immunoexpression of Ezrin and CD44 in patients with osteosarcoma. Boldrini E, et al. J Pediatr Hematol Oncol, 2010 Aug. PMID 20562647.

Ezrin immunohistochemical expression in chondrosarcomas, osteosarcomas and Ewing sarcoma family of tumors. Machado I, et al. Virchows Arch, 2010 Jul. PMID 20552365.