

Histone H3.1 Antibody (S10)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AE1016c**Specification**

Histone H3.1 Antibody (S10) - Product Information

Application	WB, IF
Primary Accession	P68431
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Concentration	1mg/ml
Isotype	Rabbit IgG
Calculated MW	15404

Histone H3.1 Antibody (S10) - Additional Information**Gene ID** 8350;8351;8352;8353;8354;8355;8356;8357;8358;8968**Other Names**

Histone H31, Histone H3/a, Histone H3/b, Histone H3/c, Histone H3/d, Histone H3/f, Histone H3/h, Histone H3/i, Histone H3/j, Histone H3/k, Histone H3/l, HIST1H3A, H3FA

Target/Specificity

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Dilution

WB~~1:500~1:1000

IF~~1:100~200

Format

affinity Purified IgG, in PBS, 0.02% sodium azide and 50% glycerol.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Histone H3.1 Antibody (S10) is for research use only and not for use in diagnostic or therapeutic procedures.

Histone H3.1 Antibody (S10) - Protein Information**Name** H3C1 ([HGNC:4766](#))**Synonyms** H3FA, HIST1H3A

Function

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Cellular Location


Nucleus. Chromosome.

Histone H3.1 Antibody (S10) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Histone H3.1 Antibody (S10) - Images



Western blot analysis of extract from HeLa cells using Histone H3.1 Antibody (S10) (#AE1016c, Lane 1 and 2) and Phospho-Histone H3.1-S10 Antibody (#AE1016d, Lane 3 and 4).



Immunofluorescence staining of methanol-fixed HeLa cells using Histone H3.1 Antibody (S10) (#AE1016c, Red).

Histone H3.1 Antibody (S10) - Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.

Histone H3.1 Antibody (S10) - References

PHF8 activates transcription of rRNA genes through H3K4me3 binding and H3K9me1/2 demethylation. Feng W, et al. Nat Struct Mol Biol, 2010 Apr. PMID 20208542.
Nucleosome formation activity of human somatic nuclear autoantigenic sperm protein (sNASP). Osakabe A, et al. J Biol Chem, 2010 Apr 16. PMID 20167597.
Structural biology of human H3K9 methyltransferases. Wu H, et al. PLoS One, 2010 Jan 11. PMID 20084102.
Molecular functions of the histone acetyltransferase chaperone complex Rtt109-Vps75. Berndsen CE, et al. Nat Struct Mol Biol, 2008 Sep. PMID 19172748.
Np95 is a histone-binding protein endowed with ubiquitin ligase activity. Citterio E, et al. Mol Cell Biol, 2004 Mar. PMID 14993289.