

Goat Anti-ADRA2A Antibody
Peptide-affinity purified goat antibody
Catalog # AF1031a**Specification**

Goat Anti-ADRA2A Antibody - Product Information

Application	IF, ICC, E
Primary Accession	P08913
Other Accession	NP_000672 , 150
Reactivity	Mouse
Predicted	Human, Rat, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	50647

Goat Anti-ADRA2A Antibody - Additional Information**Gene ID** 150**Other Names**

Alpha-2A adrenergic receptor, Alpha-2 adrenergic receptor subtype C10, Alpha-2A adrenoreceptor, Alpha-2A adrenoceptor, Alpha-2AAR, ADRA2A, ADRA2R, ADRAR

Dilution

IF~~1:50~200

ICC~~N/A

E~~N/A

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-ADRA2A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-ADRA2A Antibody - Protein Information**Name** ADRA2A ([HGNC:281](#))**Synonyms** ADRA2R, ADRAR

Function

Alpha-2 adrenergic receptors mediate the catecholamine- induced inhibition of adenylate cyclase through the action of G proteins. The rank order of potency for agonists of this receptor is oxymetazoline > clonidine > epinephrine > norepinephrine > phenylephrine > dopamine > p-synephrine > p-tyramine > serotonin = p- octopamine. For antagonists, the rank order is yohimbine > phentolamine = mianserine > chlorpromazine = spiperone = prazosin > propranolol > alprenolol = pindolol.

Cellular Location

Cell membrane; Multi-pass membrane protein

Goat Anti-ADRA2A Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-ADRA2A Antibody - Images**Goat Anti-ADRA2A Antibody - Background**

Alpha-2-adrenergic receptors are members of the G protein-coupled receptor superfamily. They include 3 highly homologous subtypes: alpha2A, alpha2B, and alpha2C. These receptors have a critical role in regulating neurotransmitter release from sympathetic nerves and from adrenergic neurons in the central nervous system. Studies in mouse revealed that both the alpha2A and alpha2C subtypes were required for normal presynaptic control of transmitter release from sympathetic nerves in the heart and from central noradrenergic neurons; the alpha2A subtype inhibited transmitter release at high stimulation frequencies, whereas the alpha2C subtype modulated neurotransmission at lower levels of nerve activity. This gene encodes alpha2A subtype and it contains no introns in either its coding or untranslated sequences.

Goat Anti-ADRA2A Antibody - References

Regional differences in cerebral perfusion associated with the alpha-2A-adrenergic receptor genotypes in attention deficit hyperactivity disorder. Kim BN, et al. J Psychiatry Neurosci, 2010 Sep. PMID 20731965.

Association between Genetic Polymorphisms of Adrenergic Receptor and Diurnal Intraocular Pressure in Japanese Normal-Tension Glaucoma. Gao Y, et al. Ophthalmology, 2010 Aug 10. PMID 20705341.

Insertion/deletion genotype of α 2B-adrenergic receptor gene polymorphism is associated with silent myocardial ischemia in patients with type 2 diabetes mellitus. Chen QJ, et al. Clin Biochem, 2010 Oct. PMID 20692245.

Variation at the NFATC2 Locus Increases the Risk of Thiazolidinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.

Resting sympathetic nerve activity is related to age, sex and arterial pressure but not to α 2-adrenergic receptor subtype. Maqbool A, et al. J Hypertens, 2010 Oct. PMID 20613626.